CUMULATIVE AUTHOR INDEX
Journal of Medical Entomology
Volume 40, 2003

A

Abdeen, Z. A., 321
Abdoulaye, D., 195
Acosta, M., 750
Adak, T., 58
Adams, T., 849
Agui, N., 468
Akida, J., 706
Aksoy, S., 755
Alexander, B., 615
Allan, S. A., 849
Almazán Garcia, C., 942
Alten, B., 930
Alves, F. P., 636
Alzogaray, R. A., 447
Amaku, M., 170
Amaral, J. C. O. F., 379
Anand, S., 534
Anderson, J. F., 528
Andreadis, T. G., 512, 528
Antonelli, A. L., 159
Apperson, C. S., 841
Arlian, L. G., 508
Arnbruster, P., 356
Arrivillaga, J., 615
Aum-Aung, B., 876
Ayres, C. F. J., 430

B

Bacon, R. M., 590
Bai, Y., 329
Baimai, V., 1
Baird, C. R., 159
Balkew, M., 141
Bangs, M. J., 876
Barbosa, S. E., 411
Barker, C. M., 403
Barnard, D. R., 463, 653
Barnett, S., 678
Barros-Battesti, D. M., 116
Beaty, B. J., 199
Becnel, J. J., 512
Beier, J. C., 657, 777, 921
Bennett, R. G., 159
Bernier, U. R., 463, 653
Biggins, D. E., 718
Birungi, J., 356
Bødker, R., 706
Boonchu, N., 983
Booth, M. M., 653
Boothby, J. T., 534
Bowers, D. F., 698
Bradley, T. J., 903
Braks, M. A. H., 785
Brantly, E., 921
Braverman, Y., 518
Bretas, G., 921
Brown, D. T., 698
Buczek, A., 690
Bunnell, J. E., 570
Buonaccorsi, J. P., 6
Burge, B., 455
Burkett, D. A., 580
Burkhalter, K., 95
Burkhalter, K. L., 361
Bustamante, D. M., 800

C

Caglar, S. S., 930
Calvo, E., 991
Camargo, E. P., 636
Camargo, L. M. A., 636
Campbell, C. L., 957
Cantrell, S., 403
Carlton, C. E., 338
Caroci, A. S., 991
Carrara, A. S., 522
Carroll, J. F., 238, 732
Castañera, P., 475
Cazorla, D., 750
Chaiwong, T., 983
Chandre, C., 195
Chandre, F., 491, 985
Chapman, H. F., 165
Chareonviriyaphap, T., 876
Chastel, C., 518
Chiles, R. E., 206, 300, 968
Chizov-Ginzburg, A., 518
Cho, H.-W., 580
Chomsky, M. S., 555
Chu, M. C., 329
Clark, D. C., 100
Clark, J. M., 628, 795
Clarke, F., 245
Claveria, F. G., 770
Coetzee, M., 36, 664
Coleman, C. G., 698
Coleman, R. E., 64, 455, 580
Collins, J. K., 577
Combe, C., 436
Conn, J. E., 356, 379
Cook, J. A., 682
Cooper, R. D., 68
Corbel, V., 985
Cornel, A. J., 36
Cornet, J.-P., 230
Cruz-Celis, A., 912
Cunha, R. P. A., 636
Cushing, P. E., 159

D

Damsky, W. E., Jr., 356
Das, A., 570
Da Silva, A. N. M., 379
da Silva, M. J. N., 116
Davé, K., 95
Davey, R. B., 482
Davidson, I., 518
Davis, B. S., 403
Day, J. F., 253, 547
Debboun, M., 64, 68, 293
De Bianchi, A. G., 991
Demay, D. J., 361
Denicola, A. J., 642
de Oliveira, M. F., 116
Desch, C. E., 146
De Souza, R. T. L., 379
Dia, I., 279
Dias-Lima, A. G., 989
Dick, C. W., 813
Diop, T., 279
Diotaiuti, L., 411
Dohm, D. J., 580
Dolan, M. C., 311
Dorn, P. L., 436
Dos Santos, C. C. B., 379
Drolet, B. S., 957
Duarte, J. M. B., 268
Dujardin, J. P., 411
Duque-De-Melo, M., 411
Durden, L. A., 91, 682, 890
Durlacher, R. R., 636

E

Edman, J. D., 6, 628, 795
Egan, D., 723
Eisen, L., 311
Elias, S., 179
Emmerich, E., 95
Enriquez, M. E., 800
Enscore, R. E., 718
Espinosa, L., 912
Estrada-Franco, J. G., 306
Estrada-Peña, A., 766
Etang, J., 491
Evangelista, L. G., 133

F

Fang, Q. Q., 890
Fang, Y., 206, 300, 968
Faulde, M. K., 935
Fekadu, H., 921
Feliciangeli, M. D., 615
Fernández, B., 95
Fernández, Z., 522

CUMULATIVE AUTHOR INDEX
Journal of Medical Entomology
Volume 40, 2003

A

Abdeen, Z. A., 321
Abdoulaye, D., 195
Acosta, M., 750
Adak, T., 58
Adams, T., 849
Agui, N., 468
Akida, J., 706
Aksoy, S., 755
Alexander, B., 615
Allan, S. A., 849
Almazán Garcia, C., 942
Alten, B., 930
Alves, F. P., 636
Alzogaray, R. A., 447
Amaku, M., 170
Amaral, J. C. O. F., 379
Anand, S., 534
Anderson, J. F., 528
Andreadis, T. G., 512, 528
Antonelli, A. L., 159
Apperson, C. S., 841
Arlian, L. G., 508
Arnbruster, P., 356
Arrivillaga, J., 615
Aum-Aung, B., 876
Ayres, C. F. J., 430

B

Bacon, R. M., 590
Bai, Y., 329
Baimai, V., 1
Baird, C. R., 159
Balkew, M., 141
Bangs, M. J., 876
Barbosa, S. E., 411
Barker, C. M., 403
Barnard, D. R., 463, 653
Barnett, S., 678
Barros-Battesti, D. M., 116
Beaty, B. J., 199
Becnel, J. J., 512
Beier, J. C., 657, 777, 921
Bennett, R. G., 159
Bernier, U. R., 463, 653
Biggins, D. E., 718
Birungi, J., 356
Bødker, R., 706
Boonchu, N., 983
Booth, M. M., 653
Boothby, J. T., 534
Bowers, D. F., 698
Bradley, T. J., 903
Braks, M. A. H., 785
Brantly, E., 921
Braverman, Y., 518
Bretas, G., 921
Brown, D. T., 698
Buczek, A., 690
Bunnell, J. E., 570
Buonaccorsi, J. P., 6
Burge, B., 455
Burkett, D. A., 580
Burkhalter, K., 95
Burkhalter, K. L., 361
Bustamante, D. M., 800

C

Caglar, S. S., 930
Calvo, E., 991
Camargo, E. P., 636
Camargo, L. M. A., 636
Campbell, C. L., 957
Cantrell, S., 403
Carlton, C. E., 338
Caroci, A. S., 991
Carrara, A. S., 522
Carroll, J. F., 238, 732
Castañera, P., 475
Cazorla, D., 750
Chaiwong, T., 983
Chandre, C., 195
Chandre, F., 491, 985
Chapman, H. F., 165
Chareonviriyaphap, T., 876
Chastel, C., 518
Chiles, R. E., 206, 300, 968
Chizov-Ginzburg, A., 518
Cho, H.-W., 580
Chomsky, M. S., 555
Chu, M. C., 329
Clark, D. C., 100
Clark, J. M., 628, 795
Clarke, F., 245
Claveria, F. G., 770
Coetzee, M., 36, 664
Coleman, C. G., 698
Coleman, R. E., 64, 455, 580
Collins, J. K., 577
Combe, C., 436
Conn, J. E., 356, 379
Cook, J. A., 682
Cooper, R. D., 68
Corbel, V., 985
Cornel, A. J., 36
Cornet, J.-P., 230
Cruz-Celis, A., 912
Cunha, R. P. A., 636
Cushing, P. E., 159

D

Damsky, W. E., Jr., 356
Das, A., 570
Da Silva, A. N. M., 379
da Silva, M. J. N., 116
Davé, K., 95
Davey, R. B., 482
Davidson, I., 518
Davis, B. S., 403
Day, J. F., 253, 547
Debboun, M., 64, 68, 293
De Bianchi, A. G., 991
Demay, D. J., 361
Denicola, A. J., 642
de Oliveira, M. F., 116
Desch, C. E., 146
De Souza, R. T. L., 379
Dia, I., 279
Dias-Lima, A. G., 989
Dick, C. W., 813
Diop, T., 279
Diotaiuti, L., 411
Dohm, D. J., 580
Dolan, M. C., 311
Dorn, P. L., 436
Dos Santos, C. C. B., 379
Drolet, B. S., 957
Duarte, J. M. B., 268
Dujardin, J. P., 411
Duque-De-Melo, M., 411
Durden, L. A., 91, 682, 890
Durlacher, R. R., 636

E

Edman, J. D., 6, 628, 795
Egan, D., 723
Eisen, L., 311
Elias, S., 179
Emmerich, E., 95
Enriquez, M. E., 800
Enscore, R. E., 718
Espinosa, L., 912
Estrada-Franco, J. G., 306
Estrada-Peña, A., 766
Etang, J., 491
Evangelista, L. G., 133

F

Fang, Q. Q., 890
Fang, Y., 206, 300, 968
Faulde, M. K., 935
Fekadu, H., 921
Feliciangeli, M. D., 615
Fernández, B., 95
Fernández, Z., 522

Fernando, K., 91
Ferreira, F., 170
Fettene, M., 112
Fondjo, E., 491
Fontenille, D., 279, 491
Forattini, O. P., 522
Frances, S. P., 68
Frederic, S., 195
Fujisaki, K., 770
Furtado, A. F., 430
Fussell, E. M., 361

G

Gage, K. L., 329, 718
Galiza, D., 379
Gannon, M. R., 813
García-Prieto, L., 996
Garruto, R. M., 119
Gebre-Michael, T., 141
Geher, K. B., 540
Gemeno, C., 841
George, J. E., 482, 942
Ghebregziabiher, A., 921
Ghebremeskel, T., 921
Gil, L. H. S., 636
Gilmore, R. D., Jr., 590
Gimnig, J. E., 421
Ginter, D. L., 275
Giordano, R., 356
Githeko, A. K., 777
Githure, J., 921
Githure, J. I., 657, 777
Glass, G. E., 570
Goddard, J., 686
Goddard, L. B., 743
Godsey, M. S., 361
Goldfarb, L., 119
Gonzalez, J.-P., 230
González-Acuña, D., 766
González Audino, P., 447
Gonzalez-Ceron, L., 371
Gottfried, K., 95
Gouagna, L. C., 657
Grant, A. J., 284
Gray, E. M., 903
Gray, J., 723
Green, E. N., 206, 300, 968
Gu, W., 595
Guerrero, F. D., 942
Guglielmone, A. A., 766
Guillet, P., 491
Gunter, J. T., 777
Gupta, R. K., 68
Gutierrez, A., 436
Guzmán-Cornejo, C., 996

H

Hall, R. A., 82, 249, 950
Hall, R. D., 559
Hanna, A. A., 73
Harrington, L. C., 6
Haskell, N. H., 559
Hawley, W. A., 421
Hernandez-Avila, J. E., 371
Higgs, S., 199
Hightower, A., 421
Hillier, A., 146
Hinkle, N. C., 678
Hirunkanokpun, S., 230
Ho, P., 498
Hoffmann, E. J., 607
Hofman, T., 690
Holden, K., 534
Holman, M. S., 179
Honório, N. A., 785
Hougard, J.-M., 985
Hribar, L. J., 361
Hsu, M.-H., 123
Hughes, J. M., 165
Hugo, L. E., 897
Humber, R. A., 451
Hung, R. W., 555
Hunt, A., 95
Hunt, R. H., 664

I

Igarashi, I., 770

J

Jacob, B. G., 777
Jacobs, M. B., 964
Jaenson, T. G. T., 375
James, S. S., 361
Janet, H., 195
Jensenius, M., 245
Jiménez, M.-L., 18
Johansen, C. A., 249
Johnson, B. J. B., 590
Johnson, J., 95
Johnston, A. N., 275
Jones, J. W., 455
Jordan, R. A., 395, 555
Joy, J. E., 73
Juliano, S. A., 785

K

Kabiru, E. W., 657
Kadota, K., 770
Kalra, N. L., 747
Kamau, L., 664
Kasai, S., 468
Kasumba, I., 755
Kaufman, M., 841
Kay, B. H., 82, 165, 897, 950
Kaynas, S., 930
Keating, J., 777
Keirans, J. E., 116, 890
Kengne, P., 279
Kennedy, B. A., 73
Kent, R. J., 223
Kerst, A., 95
Khanfar, H. M., 321
Khlaimanee, N., 64
Khrimian, A., 293
Killeen, G. F., 657
Kilpatrick, H. J., 642
Kim, H.-C., 580
Kisinza, W., 706
Kiszewski, A., 860
Kittayapong, P., 1, 230
Klein, T. A., 580
Kline, D. L., 284, 463, 653
Klun, J. A., 293
Knight, J. W., 882
Knols, B. G. J., 657
Knox, T. B., 950
Koekemoer, L. L., 664
Kollars, T. M., Jr., 103
Kosoy, M. Y., 329
Kott, S., 436
Krafsur, E. S., 30
Kramer, L. D., 215
Kramer, M., 238, 293
Krantz, G. W., 593
Kuntalue, B., 259
Kurahashi, H., 983

L

Labruna, M. B., 116, 170, 268
Lacer, L. D., 223
Lacerda, R. N. L., 379
Lacombe, E. H., 179
Lampman, R., 595
Lane, R. S., 311
Lang, J. D., 387
Langley, P., 755
Lanier, W. T., 159
Lanzaro, G. C., 615
Laver, T., 206
Lavi, M., 585
Lavigne, G. R., 179
Lee, W.-J., 580
Leepitakrat, W., 64
Lehmann, T., 421
Leite, A. C. R., 133
Lerdthusnee, K., 64
Lertthamnongtham, S., 259
Letchworth, G. J., 957
Lewis, D. J., 830
Li, A. Y., 482, 942
Licht, M., 421
Limesand, K. H., 199
Linardi, P. M., 411
Lindsay, S. W., 706
Lindström, A., 375
Little, W. E., 813
Llinas, J., 18
Lockwood, J. A., 957
Long, S. W., 1000
Lopez, R., 912
Lord, C. C., 253
Lothrop, B., 820
Lothrop, H. D., 820
Lounibos, L. P., 785
Lourenço-De-Oliveira, R., 785
Lowell, J. L., 329

Lubelczyk, C., 179
Luz, C., 451
Lysyk, T. J., 22, 185

M
MacKenzie, J. S., 82, 249
Mahapibul, P., 455
Maharaj, R., 737
Mahmood, F., 206, 300
Main, A. J., 528
Majak, W., 185
Maloney, F. A., Jr., 108
Manga, L., 491
Mann, R. B., 91
Marc, H. J., 195
Margaryan, A., 293
Margonari, C., 411
Marinotti, O., 991
Markowski, D., 555
Marquez, J. G., 30
Marquine, M., 865
Martinez, V. M., 206, 968
Massung, R. F., 534
Matsuoka, H., 441
Mbogo, C., 921
Mbogo, C. M., 777
McAbee, R. D., 36
McHolland, L. E., 348
Mclaughlin, J., 849
McNemee, R. B., Jr., 108
Mecham, J. O., 348
Meisch, M. V., 223
Melgar, S., 436
Melo-Santos, M. A. V., 430
Mendez, D., 371
Menezes, D., 989
Mengistu, S., 921
Meola, S., 755
Mercer, D. R., 807
Metzner, J. A., 170
Michalik, J., 690
Mihara, M., 468
Miller, J. R., 607
Miller, R. J., 482
Mimpfoundi, R., 491
Mogi, M., 150
Moncayo, A. C., 522
Monkanna, T., 64
Monroy, C. M., 436
Monroy, M. C., 800
Montenieri, J. A., 329, 718
Montgomery, B. L., 249
Morales-Malacara, J. B., 996
Mores, C. N., 580
Moribayashi, A., 441
Morshed, M. G., 91
Mougabure Cueto, G., 447
Msangeni, H. A., 706
Mulla, M. S., 672
Mullens, B. A., 352, 628, 795
Mumcuoglu, K. Y., 585
Mungviriya, S., 64
Munstermann, L. E., 356, 882
Munyekenye, G. O., 664
Murrell, B. P., 682
Mutebi, J.-P., 615

N
Nagpal, B. N., 747
Nakajima, Y., 78
Nanda, N., 58
Nasci, R., 95
Navarro, R., 306
Nayar, J. K., 882
Ngo, K. A., 215
Nisbet, D. J., 82
Nishiura, J. T., 498
Noigamol, C., 455
Noireau, F., 989
Norris, D., 615
Norris, S. L. W., 455
Northill, J. A., 249
Novak, R., 921
Novak, R. J., 595
Nunamaker, R. A., 957
N'Guessan, R., 985

O
Oda, T., 441
Ogihara, K., 78
O'Guinn, M. L., 306, 580
Okech, B. A., 657
Oliver, J. H., Jr., 103
Olson, J. K., 259
O'Neill, S. L., 1
Ordoñez, R., 912
Ostfeld, R. S., 540

P
Palacios, C., 18
Pant, C. S., 58
Panzera, F., 411
Pasteur, N., 865
Patrick, M. J., 813
Paulson, S. L., 403
Pearson, L. D., 199
Pedersen, E. M., 706
Pereira, M. C., 268
Pereira, M. H., 411
Pereira-Da-Silva, L. H., 636
Pérez-Ponce De León, G., 996
Perich, M. J., 455, 930
Perotti, M. A., 22
Philipp, M. T., 964
Piangjai, S., 983
Pichon, B., 723
Picollo, M. I., 447
Pierre, G., 195
Pierre, K., 195
Piesman, J., 311, 590
Pimenta, P. F. P., 411
Piñango, H., 615
Pinter, A., 170
Pires, H. H. R., 411
Policastro, P. F., 364
Pollack, R. J., 860
Posey, K. H., 463, 653
Póvoa, M. M., 379
Prabaripai, A., 876
Pretorius, A.-M., 245
Price, S. D., 570
Puelle, R. S., 555
Purcell, J. E., 964
Pyke, A. T., 249

Q
Quintana, M., 590

R
Rabe, M. J., 577
Radovsky, F. J., 593
Rakotoarivony, I., 279
Ramberg, F., 577
Ramsey, J. M., 912
Rand, P. W., 179
Rasgon, J., 36
Rasgon, J. L., 125
Rau, M. E., 830
Ray, K., 498
Raymond, M., 865
Regens, J. L., 777
Reisen, W. K., 206, 300, 743, 820, 968
Ribolla, P. E., 991
Ringertz, S. H., 245
Ritchie, S. A., 82, 165, 249
Robbins, R. G., 935
Robert, G. T., 195
Roch, D. K., 195
Rocha, L. F. N., 451
Rodas, A., 436
Rodas, A. G., 800
Rodriguez, M. H., 371
Roe, A. H., 159
Rogers, M., 723
Romanha, A. J., 411
Romero, G., 306
Rosa, E. P. S., 379
Rosales, R., 436
Rosales, R. G., 800
Rosenstock, S. S., 577
Roth, A. E., 743
Rouzier, V., 436
Royce, L. A., 159
Ruang-Areerate, T., 1
Ruble, R. P., 1000
Rust, M. K., 678
Rutledge, C. R., 253
Ryan, J., 95
Ryan, P. A., 897, 950

S
Salcedo, J. M. V., 636
Salgado, L., 912
Salvia, D., 436
Sames, W. J., IV, 108

Sánchez-Ramos, I., 475
Santillan, F., 371
Sawalha, S. S., 321
Schaffner, F., 865
Schal, C., 841
Schell, S. P., 957
Schleich, S. S., 455
Schlichting, C. D., 379
Schmidt, J. O., 275
Schoenly, K., 559
Schofield, C. J., 912
Schulze, T. L., 395, 555
Schwab, A. E., 830
Schwan, T. G., 364
Scott, J. D., 91
Scott, T. W., 36, 125, 743
Seery, D. B., 718
Segura, M. N. O., 379
Seulu, F., 921
Shahid, S. A., 559
Shaman, J., 547
Shayo, D., 706
Sherlock, I., 989
Shields, T. M., 570
Shililu, J., 921
Shtayeh, M. S., 321
Sikora, B., 690
Simsek, F. M., 930
Sintasath, D., 921
Sithiprasasna, R., 455
Sittertz-Bhatkar, H., 755
Skoracki, M., 690
Smith, G. A., 249
Smith, H., 352
Smith, J. M., 361
Smith, R. P., Jr, 179
Soares, R. P. P., 411
Solé-Cava, A. M., 430
Sonenshine, D. E., 849
Spangler, H. G., 275
Spielman, A., 860
Spinelli, G. R., 807
Spoto, M. C., 361
Srivastava, A., 747
Srivastava, H. C., 58
Stafford, K. C., III, 642
Stanich, M. A., 36
Stansfield, D., 678
Stark, L. M., 253, 361
Stegall-Faulk, T., 100
Stevenson, H. L., 329
Stiebel, G. D., 585
Stieglitz, M., 547
Stith, C. E., 957
Stoner, R. L., 361
Styer, L. M., 125
Subbarao, S. K., 58, 747
Sukontason, K., 259, 983
Sukontason, K. L., 259, 983
Sunahara, T., 150
Swalm, C. M., 777
Szabó, M. P. J., 268

T

Tabachnick, W. J., 253, 597
Tada, M. S., 636
Takahashi, M., 468
Takano-Lee, M., 628, 795
Tanda, D. T., 718
Tarshis, M., 585
Taylor, D., 78, 770
Teel, P., 1000
Temu, E. A., 112
Tesh, R. B., 807
Thapar, B. R., 58
Thierry, B., 195
Thijuk, N., 259
Tomita, T., 468
Toto, J.-C., 491
Townsend, A. K., 540
Trexler, J. D., 841
Tseng, M., 588
Turell, M., 95
Turell, M. J., 306, 580

U

Uchida, K., 441
Uedelhoven, W. M., 935
Uspensky, I., 119

V

van den Hurk, A. F., 82, 249
Vassena, C. V., 447
Velten, R. K., 795
Venzal, J. M., 766
Vetter, R. S., 159
Villarino, M. A., 52
Vlach, J. J., 361
Vogtsberger, R. C., 259
Vossbrinck, C. R., 528
Vulule, J. M., 421
Vyszenski-Moher, D. L., 508

W

Waggoner, M. M., 678
Waghela, S. D., 52
Wagner, G. G., 52
Wahid, I., 150
Walker, E., 841
Wallace, L., 841
Walter, S., 770
Warburg, A., 321
Wasieloski, L. P., Jr., 580
Watson, D. W., 841
Watson, E. J., 338
Watts, D. M., 807
Weaver, S. C., 522
Webster, F. X., 849
White, D. M., 957
White, S. E., 512
Wikel, S. K., 528
Wilson, W. C., 957
Wirtz, R., 95
Wirtz, R. A., 379
Wright, S. M., 100

X

Xu, G., 890

Y

Yadav, R. S., 58
Yaguchi, N., 468
Yamakawa, M., 78
Yan, G., 657
Ye-Ebiyo, Y., 860
Yoon, K. S., 628
Yost, R. A., 653
Yu, X.-J., 1000

Z

Zahiri, N. S., 672
Zaim, M., 58
Zeichner, B. C., 455
Zerba, E. N., 447
Zerom, M., 921
Zhang, J., 1000
Zhang, W., 559
Zhang, X., 1000
Zias, J., 585
Zieler, H., 636
Zurek, L., 841

CUMULATIVE SUBJECT INDEX
Journal of Medical Entomology
Volume 40, 2003

A

abundance, 352, 785
  Brazil, 785
  Culicoides, 352
  dengue vectors, 785
  midges, 352
  occurrence, 785
  parity, 352
  United States, 785
Acari, 146, 996
  bats, 996
  Demodecidae, 146
  domestic dog, 146
  ectoparasite, 146
  infracommunities, 996
  Mexico, 996
  mite, 146
  Tadarida brasiliensis mexicana, 996
acaricide resistance,
  Boophilus microplus, 482
  cattle tick, 482
  coumaphos, 482
  diazinon, 482
  organophosphate, 482
acetylcholinesterase,
  cDNA, 890
  Dermacentor variabilis, 890
  Rhipicephalus sanguineus, 890
  ticks, 890
adults,
  Chagas' disease, 18
  habitat, 18
  kissing bugs, 18
  nymphs, 18
Aedes,
  artificial feeding, 588
  Parafilm M, 588
Aedes aegypti, 6, 293, 830, 950
  (1S, 2′S)-2-methylpiperidinyl-3-cyclohexen-1-carboxamide, 293
  2-(2-hydroxyethyl)-1-piperidinecarboxylic acid 1-methylpropyl ester, 293
  Anopheles stephensi, 293
  biological control, 830
  Bootstrapping, 6
  dengue, 950
  mark-release, 6
  N,N-diethyl-3-methylbenzamide, 293
  nonlinear regression, 6
  oviposition, 830
  Plagiorchis elegans, 830
  population ecology, 830
  statistical inference, 6
  survival rates, 6
  transmission, 950
  vector competence, 950
  virus, 950
Aedes albopictus, 73, 356, 403, 522, 698, 841
  arbovirus, 403
  attractants, 841
  bacteria, 841
  electroantennogram, 841
  encephalitis, 403
  La Crosse virus, 403
  longevity, 522
  midgut visceral muscles, 698
  mitochondrial DNA, 356
  mosquito larval habitats, 73
  mosquito pathology, 698
  mosquito strains, 522
  Ochlerotatus triseriatus, 73, 403
  organic infusion, 841
  oviposition, 403, 841
  salivary glands, 698
  Sindbis virus, 698
  vector competence, 522
  Venezuelan equine encephalitis, 522
  West Virginia, 73
  Wolbachia, 356
  wsp gene, 356
Aedes mosquitoes,
  control, 455
  dengue, 455
  lethal ovitrap, 455
  surveillance, 455
Aedes vexans,
  Anopheles walkeri, 607
  carbon dioxide, 607
  Coquillettidae perturbans, 607
  orientation, 607
Africanized honey bees,
  Apis mellifera, 275
  attacks, 275
  citronella, 275
  insect repellents, 275
  NN-diethyl-*m*-toluamide, 275
age-structure,
  Culex pipiens, 125
  popcorn, 125
  vector-borne disease, 125
  Wolbachia, 125
Alaska,
  Ixodes angustus, 682
  Leptinus occidentamericanus, 682
  mammalian hosts, 682
allozymes,
  Anopheles claviger, 865
  Europe, 865
  population genetics, 865
  sibling species, 865
Amazon Region,
  Anopheles, 636
  malaria, 636
  Plasmodium, 636
  transmission, 636
Amblyomma,
  16S rRNA, 590
  Borrelia, 590
  flagellin, 590
  Missouri, 590
Amblyomma americanum, 395, 528, 686, 732
  blacklegged tick, 732
  Dermacentor andersoni, 528
  Dermacentor variabilis, 528
  disease transmission, 686
  infection, 686
  Ixodes scapularis, 395, 528, 732
  lone star tick, 732
  questing, 395
  Rickettsia parkeri, 686
  weather, 395
  West Nile virus, 528
Amblyomma cajennense, 170, 268
  Amblyomma triste, 268
  Anocentor nitens, 268
  Blastocerus dichotomus, 268
  Boophilus microplus, 268
  Brazil, 170
  diapause, 170
  marsh deer, 268
  premolt period, 170
Amblyomma hebraeum,
  DEET, 245
  KBR 3023, 245
  piperidine, 245
  repellents, 245
  tick, 245
Amblyomma testudinarium,
  Haemaphysalis ornithophila, 230
  rickettsiae, 230
  Thailand, 230
  tick, 230

Amblyomma triste,
Amblyomma cajennense, 268
Anocentor nitens, 268
Blastocerus dichotomus, 268
Boophilus microplus, 268
marsh deer, 268
americanum,
B. lonestari, 100
lyme-like, 100
STARI, 100
amino acids,
Anopheles stephensi, 441
infusion, 441
mating, 441
oogenesis, 441
amplification,
Culex nigripalpus, 547
hydrology, 547
St. Louis encephalitis, 547
wild birds, 547
Anaplasma phagocytophilum,
Borrelia burgdorferi, 534
California, 534
Ehrlichia chaffeensis, 534
Ixodes pacificus, 534
Anocentor nitens,
Amblyomma cajennense, 268
Amblyomma triste, 268
Blastocerus dichotomus, 268
Boophilus microplus, 268
marsh deer, 268
Anopheles, 636, 991
Amazon Region, 636
malaria, 636
malaria vector, 991
Plasmodium, 636
transmission, 636
trypsin, 991
Anopheles albimanus,
bacteria, 371
midgut, 371
Plasmodium vivax, 371
sporogony, 371
Anopheles aquasalis,
Anopheles darlingi, 379
malaria incidence, 379
malaria vector, 379
northeastern Brazil, 379
Anopheles arabiensis,
life tables, 737
seasonal conditions, 737
Anopheles atropos,
Deinocerites cancer, 361
Florida Keys, 361
Ochlerotatus taeniorhynchus, 361
West Nile virus, 361
Anopheles claviger,
allozymes, 865
Europe, 865
population genetics, 865
sibling species, 865
Anopheles culicifacies,
bifenthrin, 58
excito-repellency, 58
malathion, 58
spraying, 58
vector density, 58
Anopheles darlingi,
Anopheles aquasalis, 379
malaria incidence, 379
malaria vector, 379
northeastern Brazil, 379
Anopheles dispersal,
genetic-relatedness malaria, 421
kinship, 421
population structure, 421
Anopheles farauti,
bednet, 68
malaria, 68
Papua New Guinea, 68
permethrin, 68
personal protection, 68
Anopheles funestus,
chromosomal inversions, 664
malaria, 664
species composition, 664
variability, 664
Anopheles gambiae, 112, 195, 491, 985
Anopheles quadriannulatus sp. B, 112
Cameroon, 491
Côte d'Ivoire, 985
Ethiopia, 112
insecticide combinations, 985
insecticide resistance, 491
insecticide-treated nets, 985
insensitive acetylcholinesterase, 985
kdr mutation, 195
molecular M form, 195
PCR assay, 112
tropical savannah area, 195
Anopheles gambiae s.l. complex,
geographic information systems (GIS), 777
land use land cover change (LULC), 777
multispectral thermal imager (MTI), 777
Anopheles larval habitats,
Eritrea, 921
larval control, 921
spatial temporal distribution, 921
Anopheles minimus,
blood feeding, 876
exophagy, 876
mosquito abundance, 876
Thailand, 876
Anopheles quadriannulatus sp. B,
Anopheles gambiae, 112
Ethiopia, 112
PCR assay, 112
Anopheles spp.,
highland malaria, 706
malaria transmission, 706
Tanzania, 706
vectorial capacity, 706
Anopheles stephensi, 293, 441
(1*S*, 2′*S*)-2-methylpiperidinyl-3-cyclohexen-1-carboxamide, 293
2-(2-hydroxyethyl)-1-piperidinecarboxylic acid 1-methylpropyl ester, 293
Aedes aegypti, 293
amino acids, 441
infusion, 441
mating, 441
N,N-diethyl-3-methylbenzamide, 293
oogenesis, 441
Anopheles stephensi type form,
ecological variants, 747
spiracular index, 747
taxonomic tools, 747
thoracic spiracle length, 747
variant *mysorensis*, 747
Anopheles survival,
blood, 657
indoor climate, 657
sugar, 657
vectorial capacity, 657
Anopheles walkeri,
Aedes vexans, 607
carbon dioxide, 607
Coquillettidae perturbans, 607
orientation, 607
Anoplura,
head lice, 795
in vitro rearing, 795
Pediculidae, 795
Pediculus humanus capitis, 795
Anthropology Research Facility,
carrion-arthropod succession, 559
forensic entomology, 559
nutrient enrichment, 559
antibacterial activity,
hemoglobin fragment, 78
midgut, 78
Ornithodoros moubata, 78
antibody, 300, 968
chronic infection, 968
experimental infection, 968
house finches, 300
St. Louis encephalitis virus, 300, 968
viral persistence, 300
western equine

encephalomyelitis virus, 968
Apis mellifera,
Africanized honey bees, 275
attacks, 275
citronella, 275
insect repellents, 275
NN-diethyl-*m*-toluamide, 275
Arachnida,
non-native, 159
range expansion, 159
Tegenaria agrestis, 159
arbovirus, 95, 223, 249, 253, 403, 807
Aedes albopictus, 403
Arkansas, 223
Australia, 249
biting rates, 807
blue jay, 223
developmental substrates, 807
encephalitis, 403
field transmission, 253
host-seeking, 807
infection rate, 253
Japanese encephalitis, 249
La Crosse virus, 403
mosquito, 223
mosquito trap, 249
Ochlerotatus triseriatus, 403
Oropouche virus, 807
oviposition, 403
polymerase chain reaction, 249
rapid detection, 95
St. Louis encephalitis, 95
surveillance, 95, 223, 249
West Nile virus, 95, 223, 253
wicking assay, 95
archaeology,
body louse, 585
Israel, 585
Masada, 585
Pediculus humanus humanus, 585
archeological site,
Chagas disease, 800
invasion, 800
sylvatic habitats, 800
Triatoma dimidiata, 800
area repellent,
cis/*trans* allethrin, 930
Ochlerotatus caspius, 930
Phlebotomus papatasi, 930
Thermacell, 930
Argas (*P.*) *keiransi* n. sp.,
Chile, 766
Falconiformes, 766
Arkansas,
arbovirus, 223
blue jay, 223
mosquito, 223
surveillance, 223
West Nile virus, 223
arthropod succession,
carrion ecology, 338
forensic entomology, 338
poaching, 338
wildlife, 338
arthropods,
permethrin, 935
repellents, 935
uniforms, 935
artificial feeding, 588, 628
Aedes, 588
head lice, 628
Parafilm M, 588
Pediculidae, 628
Pediculus humanus capitis, 628
Phthiraptera, 628
attacks,
Africanized honey bees, 275
Apis mellifera, 275
citronella, 275
insect repellents, 275
NN-diethyl-*m*-toluamide, 275
attractants, 284, 841
Aedes albopictus, 841
bacteria, 841
carbon dioxide, 284
Culicoides furens, 284
electroantennogram, 841
electrophysiology, 284
olfaction, 284
organic infusion, 841
oviposition, 841
Australia, 82, 249, 897
arbovirus, 249
autogeny, 897
fecundity, 897
Japanese encephalitis, 82, 249
mosquito trap, 249
mosquitoes, 82
Ochlerotatus vigilax, 897
polymerase chain reaction, 249
surveillance, 249
vector competence, 82
wing length, 897
autogeny, 150, 897
Australia, 897
fecundity, 897
male mosquito, 150
mandible, 150
maxilla, 150
Ochlerotatus vigilax, 897
partial autogeny, 150
wing length, 897

B

B. burgdorferi,
I. ricinus, 690
reservoir, 690
small rodents, 690
urban forests, 690
B. lonestari,
americanum, 100
lyme-like, 100
STARI, 100
Bacillus sphaericus,
Bacillus thuringiensis ssp. israelensis, 672
mixture, 672
mosquitoes, 672
resistance, 672
Bacillus thuringiensis israelensis,
feeding by larval mosquitoes, 860
maize pollen, 860
phagostimulant, 860
Bacillus thuringiensis ssp. israelensis,
Bacillus sphaericus, 672
mixture, 672
mosquitoes, 672
resistance, 672
bacteria, 22, 371, 841
Aedes albopictus, 841
Anopheles albimanus, 371
attractants, 841
development, 22
electroantennogram, 841
growth, 22
Hematobia irritans, 22
larval survival, 22
media, 22
midgut, 371
organic infusion, 841
oviposition, 841
Plasmodium vivax, 371
sporogony, 371
Baculoviridae,
Culex, 512
CuniNPV, 512
infectivity, 512
mosquito, 512
Barbour-Stoenner-Kelly medium,
Lyme disease, 364
Peromyscus leucopus, 364
spirochete, 364
tick, 364
Bartonella,
black-tailed prairie dog, 329
multiplex polymerase chain reaction, 329
Oropsylla, 329
Rickettsia, 329
Yersinia pestis, 329
bat guano habitat,
Coprolactistus whitakeri, 593
Mitonyssoides stercoralis, 593
parasitism, 593
predation, 593
bats,
Acari, 996
infracommunities, 996
Mexico, 996

Tadarida brasiliensis mexicana, 996
bednet,
Anopheles farauti, 68
malaria, 68
Papua New Guinea, 68
permethrin, 68
personal protection, 68
behavioral bioassay,
kairomones, 653
mosquito attractants, 653
behavioral observations,
Capstar, 678
cat flea, 678
Ctenocephalides felis, 678
nitenpyram, 678
bifenthrin,
Anopheles culicifacies, 58
excito-repellency, 58
malathion, 58
spraying, 58
vector density, 58
bioassay,
Pediculus humanus capitis, 447
permethrin resistance, 447
biological control,
Aedes aegypti, 830
oviposition, 830
Plagiorchis elegans, 830
population ecology, 830
biotopes,
Ixodes ricinus, 375
Sweden, 375
tick sampling, 375
vegetation types, 375
Birds,
Borrelia burgdorferi, 540
Ixodes, 540
Lyme disease, 540
risk factors, 540
biting activity,
leishmaniasis, 321
Phlebotomus, 321
resting sites, 321
West Bank, 321
biting rates,
arbovirus, 807
developmental substrates, 807
host-seeking, 807
Oropouche virus, 807
blacklegged tick, 91, 238, 732
Amblyomma americanum, 732
Borrelia burgdorferi, 91
Ixodes scapularis, 91, 732
lone star tick, 732
Lyme disease, 91
model, 238
Ontario, 91
Rondeau Provincial Park, 91
weather, 238
white-tailed deer, 238
'4-poster', 238
black-tailed prairie dog,
Bartonella, 329
multiplex polymerase chain reaction, 329
Oropsylla, 329
Rickettsia, 329
Yersinia pestis, 329
Blastocerus dichotomus,
Amblyomma cajennense, 268
Amblyomma triste, 268
Anocentor nitens, 268
Boophilus microplus, 268
marsh deer, 268
blood,
Anopheles survival, 657
indoor climate, 657
sugar, 657
vectorial capacity, 657
blood feeding,
Anopheles minimus, 876
exophagy, 876
mosquito abundance, 876
Thailand, 876
blood meal analysis,
Ixodes ricinus, 723
reservoir host identification, 723
tick-borne zoonoses, 723
bloodmeal,
Culex, 215
West Nile, 215
blue jay,
arbovirus, 223
Arkansas, 223
mosquito, 223
surveillance, 223
West Nile virus, 223
bluetongue,
Culicoides mohave, 577
Culicoides sonorensis, 577
epizootic hemorrhagic disease, 577
vectors, 577
bluetongue viruses,
cell lines, 348
Culicoides sonorensis, 348
epizootic hemorrhagic disease viruses, 348
field populations, 348
isozyme phenotyping, 348
body louse, 468, 585
archaeology, 585
head louse, 468
insecticide-resistance, 468
Israel, 585
Masada, 585
para-orthologous sodium channel gene, 468
pediculicide, 468
Pediculus humanus humanus, 585
Boophilus microplus, 52, 268, 482
acaricide resistance, 482
Amblyomma cajennense, 268
Amblyomma triste, 268
Anocentor nitens, 268
Blastocerus dichotomus, 268
cattle tick, 482
coumaphos, 482
diazinon, 482
esterase purification, 52
esterases, 52
insecticide resistance, 52
marsh deer, 268
organophosphate, 482
Bootstrapping,
Aedes aegypti, 6
mark-release, 6
nonlinear regression, 6
statistical inference, 6
survival rates, 6
Borrelia,
16S rRNA, 590
Amblyomma, 590
flagellin, 590
Missouri, 590
Borrelia bissettii,
feeding success, 311
Ixodes pacificus, 311
Ixodes spinipalpis, 311
rodent reservoir, 311
tick vector, 311
Borrelia burgdorferi, 91, 534, 540, 555, 964
Anaplasma phagocytophilum, 534
Birds, 540
blacklegged tick, 91
California, 534
distribution, 555
Ehrlichia chaffeensis, 534
Ixodes, 540
Ixodes pacificus, 534
Ixodes scapularis, 91, 555, 964
Louisiana, 964
Lyme borreliosis, 964
Lyme disease, 91, 540
Ontario, 91
risk factors, 540
Rondeau Provincial Park, 91
bovine,
grasshopper, 957
Melanoplus sanguinipes, 957
vesicular stomatitis virus, 957
VSV, 957
Brazil, 116, 170, 785
abundance, 785
Amblyomma cajennense, 170
dengue vectors, 785
diapause, 170
Ixodes schulzei, 116
laboratory rearing, 116
occurrence, 785

premolt period, 170
ticks, 116
United States, 785

C

California,
Anaplasma phagocytophilum, 534
Borrelia burgdorferi, 534
Ehrlichia chaffeensis, 534
Ixodes pacificus, 534
Cameroon,
Anopheles gambiae, 491
insecticide resistance, 491
Capstar,
behavioral observations, 678
cat flea, 678
Ctenocephalides felis, 678
nitenpyram, 678
carbon dioxide, 284, 607
Aedes vexans, 607
Anopheles walkeri, 607
attractants, 284
Coquillettidae perturbans, 607
Culicoides furens, 284
electrophysiology, 284
olfaction, 284
orientation, 607
carcass,
carrion, 108
decomposition, 108
Dermacentor variabilis, 108
Erethizon dorsatum, 108, 108
carpet cleaning,
Dermatophagoides, 508
Disodium octaborate tetrahydrate, 508
house dust mites, 508
carrion,
carcass, 108
decomposition, 108
Dermacentor variabilis, 108
Erethizon dorsatum, 108, 108
carrion ecology,
arthropod succession, 338
forensic entomology, 338
poaching, 338
wildlife, 338
carrion-arthropod succession,
Anthropology Research Facility, 559
forensic entomology, 559
nutrient enrichment, 559
cat flea,
behavioral observations, 678
Capstar, 678
Ctenocephalides felis, 678
nitenpyram, 678
cattle tick,
acaricide resistance, 482
Boophilus microplus, 482
coumaphos, 482
diazinon, 482
organophosphate, 482
cDNA,
acetylcholinesterase, 890
Dermacentor variabilis, 890
Rhipicephalus sanguineus, 890
ticks, 890
cell lines,
bluetongue viruses, 348
Culicoides sonorensis, 348
epizootic hemorrhagic disease viruses, 348
field populations, 348
isozyme phenotyping, 348
Chagas' disease, 18, 436, 451, 800
adults, 18
archeological site, 800
entomopathogenic fungi, 451
Evlachovaea, 451
Guatemala, 436
habitat, 18
invasion, 800
kissing bugs, 18
nymphs, 18
population genetics, 436
sylvatic habitats, 800
Triatoma dimidiata, 436, 800
Triatoma sordida, 451
vector control, 436, 451
chiggers,
Leptotrombidium, 64
personal protection, 64
repellents, 64
scrub typhus, 64
Chile,
Argas (*P.*) *keiransi* n. sp., 766
Falconiformes, 766
Chiroptera,
ectoparasites, 813
host–parasite associations, 813
Myodopsylla insignis, 813
Myotis lucifugus, 813
chromosomal inversions,
Anopheles funestus, 664
malaria, 664
species composition, 664
variability, 664
chronic infection, 206, 968
antibody, 968
cyclophosphamide, 206
dexamethasone, 206
experimental infection, 968
house finch, 206
immunosuppression, 206
mosquito blood feeding, 206
St. Louis encephalitis virus, 968
western equine encephalomyelitis virus, 968
Chrysomya rufifacies,
forensic entomology, 259
larval morphology, 259
scanning electron microscopy, 259
ultrastructure, 259
Chrysomya villeneuvi,
forensic entomology, 983
hairy maggot, 983
circadian activity,
feeding, 903
female mosquito, 903
flow-through respirometry, 903
metabolic rate, 903
cis/*trans* allethrin,
area repellent, 930
Ochlerotatus caspius, 930
Phlebotomus papatasi, 930
Thermacell, 930
citronella,
Africanized honey bees, 275
Apis mellifera, 275
attacks, 275
insect repellents, 275
NN-diethyl-*m*-toluamide, 275
cold storage,
Dermacentor andersoni, 185
laboratory colony, 185
Rocky Mountain wood tick, 185
selection, 185
tick paralysis, 185
control,
Aedes mosquitoes, 455
dengue, 455
lethal ovitrap, 455
surveillance, 455
Coprolactistus whitakeri,
bat guano habitat, 593
Mitonyssoides stercoralis, 593
parasitism, 593
predation, 593
Coquillettidae perturbans,
Aedes vexans, 607
Anopheles walkeri, 607
carbon dioxide, 607
orientation, 607
Côte d'Ivoire,
Anopheles gambiae, 985
insecticide combinations, 985
insecticide-treated nets, 985
insensitive acetylcholinesterase, 985
coumaphos,
acaricide resistance, 482
Boophilus microplus, 482
cattle tick, 482
diazinon, 482
organophosphate, 482

Ctenocephalides felis,
behavioral observations, 678
Capstar, 678
cat flea, 678
nitenpyram, 678
Culex, 215, 512
Baculoviridae, 512
bloodmeal, 215
CuniNPV, 512
infectivity, 512
mosquito, 512
West Nile, 215
Culex annulirostris,
Culex palpalis, 165
Culicidae, 165
dispersal, 165
Japanese encephalitis, 165
Culex nigripalpus,
amplification, 547
hydrology, 547
St. Louis encephalitis, 547
wild birds, 547
Culex palpalis,
Culex annulirostris, 165
Culicidae, 165
dispersal, 165
Japanese encephalitis, 165
Culex pipiens, 125, 743
age-structure, 125
Culex tarsalis, 743
popcorn, 125
vector-borne disease, 125
vertical transmission, 743
West Nile virus, 743
Wolbachia, 125
Culex pipiens quinquefasciatus,
geographic variation, 882
isozyme electrophoresis, 882
population genetics, 882
temporal variation, 882
Culex pseudes,
Honduras, 306
Mexico, 306
Ochlerotatus taeniorhynchus, 306
transmission, 306
Venezuelan equine encephalomyelitis virus, 306
Culex tarsalis, 743, 820
Culex pipiens, 743
dispersal, 820
mark-release-recapture, 820
mosquitoes, 820
vertical transmission, 743
West Nile virus, 743
Culicidae,
Culex annulirostris, 165
Culex palpalis, 165
dispersal, 165
Japanese encephalitis, 165
Culicoides,
abundance, 352
midges, 352
parity, 352
Culicoides furens,
attractants, 284
carbon dioxide, 284
electrophysiology, 284
olfaction, 284
Culicoides mohave,
bluetongue, 577
Culicoides sonorensis, 577
epizootic hemorrhagic disease, 577
vectors, 577
Culicoides sonorensis, 348, 577
bluetongue, 577
bluetongue viruses, 348
cell lines, 348
Culicoides mohave, 577
epizootic hemorrhagic disease, 577
epizootic hemorrhagic disease viruses, 348
field populations, 348
isozyme phenotyping, 348
vectors, 577
CuniNPV,
Baculoviridae, 512
Culex, 512
infectivity, 512
mosquito, 512
cyclophosphamide,
chronic infection, 206
dexamethasone, 206
house finch, 206
immunosuppression, 206
mosquito blood feeding, 206

D

Dasypodidae,
northeastern Brazil, 989
Panstrongylus lutzi, 989
silvatic ecology, 989
Triatominae, 989
decomposition,
carcass, 108
carrion, 108
Dermacentor variabilis, 108
Erethizon dorsatum, 108, 108
deer density,
Ixodes scapularis, 179
white-tailed deer, 179
deer reduction,
Ixodes scapularis, 642
Ixodiphagus hookeri, 642
parasitoid, 642
tick, 642
DEET, 245, 463
Amblyomma hebraeum, 245
dehydrolinalool, 463
inhibition, 463
KBR 3023, 245
linalool, 463
olfactometer, 463
piperidine, 245
repellents, 245
tick, 245
dehydrolinalool,
deet, 463
inhibition, 463
linalool, 463
olfactometer, 463
Deinocerites cancer,
Anopheles atropos, 361
Florida Keys, 361
Ochlerotatus taeniorhynchus, 361
West Nile virus, 361
Deltadust,
epizootic, 718
fleas, 718
insecticide, 718
plague, 718
prairie dogs, 718
Demodecidae,
Acari, 146
domestic dog, 146
ectoparasite, 146
mite, 146
dengue, 455, 950
Aedes aegypti, 950
Aedes mosquitoes, 455
control, 455
lethal ovitrap, 455
surveillance, 455
transmission, 950
vector competence, 950
virus, 950
dengue vectors,
abundance, 785
Brazil, 785
occurrence, 785
United States, 785
Dermacentor andersoni, 185, 528
Amblyomma americanum, 528
cold storage, 185
Dermacentor variabilis, 528
Ixodes scapularis, 528
laboratory colony, 185
Rocky Mountain wood tick, 185
selection, 185
tick paralysis, 185
West Nile virus, 528
Dermacentor variabilis, 108, 528, 890
acetylcholinesterase, 890
Amblyomma americanum, 528
carcass, 108
carrion, 108
cDNA, 890
decomposition, 108
Dermacentor andersoni, 528

Erethizon dorsatum, 108, 108
Ixodes scapularis, 528
Rhipicephalus sanguineus, 890
ticks, 890
West Nile virus, 528
Dermatobia hominis,
human bot fly, 133
larva, 133
midgut, 133
ultrastructure, 133
Dermatophagoides,
carpet cleaning, 508
Disodium octaborate tetrahydrate, 508
house dust mites, 508
development,
bacteria, 22
growth, 22
Hematobia irritans, 22
larval survival, 22
media, 22
developmental substrates,
arbovirus, 807
biting rates, 807
host-seeking, 807
Oropouche virus, 807
dexamethasone,
chronic infection, 206
cyclophosphamide, 206
house finch, 206
immunosuppression, 206
mosquito blood feeding, 206
diapause,
Amblyomma cajennense, 170
Brazil, 170
premolt period, 170
diazinon, 482, 942
acaricide resistance, 482
Boophilus microplus, 482
cattle tick, 482
coumaphos, 482
insecticide resistance, 942
kdr, 942
organophosphate, 482
permethrin, 942
the horn fly, 942
Diptera,
dorsal septum, 755
immunocytochemistry, 755
neuropeptides, 755
disease transmission,
Amblyomma americanum, 686
infection, 686
Rickettsia parkeri, 686
Disodium octaborate tetrahydrate,
carpet cleaning, 508
Dermatophagoides, 508
house dust mites, 508
dispersal, 165, 820
Culex annulirostris, 165
Culex palpalis, 165
Culex tarsalis, 820
Culicidae, 165
Japanese encephalitis, 165
mark-release-recapture, 820
mosquitoes, 820
distribution,
Borrelia burgdorferi, 555
Ixodes scapularis, 555
distributional range,
introduction, 119
Ixodes persulcatus, 119
northern range border, 119
reproductive range, 119
Sakha Republic (Yakutia), 119
domestic control,
pyrethroid insecticides, 912
Triatoma barberi, 912
Triatoma pallidipennis, 912
domestic dog,
Acari, 146
Demodecidae, 146
ectoparasite, 146
mite, 146
dorsal septum,
Diptera, 755
immunocytochemistry, 755
neuropeptides, 755

E

ecdysial phase,
hemocyte class, 770
Ornithodoros moubata, 770
ultrastructure, 770
ecological variants,
Anopheles stephensi type form, 747
spiracular index, 747
taxonomic tools, 747
thoracic spiracle length, 747
variant *mysorensis*, 747
ectoparasites, 146, 813
Acari, 146
Chiroptera, 813
Demodecidae, 146
domestic dog, 146
host–parasite associations, 813
mite, 146
Myodopsylla insignis, 813
Myotis lucifugus, 813
Ehrlichia chaffeensis,
Anaplasma phagocytophilum, 534
Borrelia burgdorferi, 534
California, 534
Ixodes pacificus, 534
electroantennogram,
Aedes albopictus, 841
attractants, 841
bacteria, 841
organic infusion, 841
oviposition, 841
electrophysiology,
attractants, 284
carbon dioxide, 284
Culicoides furens, 284
olfaction, 284
encephalitis,
Aedes albopictus, 403
arbovirus, 403
La Crosse virus, 403
Ochlerotatus triseriatus, 403
oviposition, 403
endosymbionts,
mosquitoes, 1
phylogeny, 1
Wolbachia, 1
wsp sequences, 1
entomopathogenic fungi,
Chagas' disease, 451
Evlachovaea, 451
Triatoma sordida, 451
vector control, 451
environmental predictors,
GIS, 570
Ixodes scapularis, 570
Poisson-normal model, 570
soil, 570
spatial correlation, 570
epizootic,
Deltadust, 718
fleas, 718
insecticide, 718
plague, 718
prairie dogs, 718
epizootic hemorrhagic disease,
bluetongue, 577
Culicoides mohave, 577
Culicoides sonorensis, 577
vectors, 577
epizootic hemorrhagic disease viruses,
bluetongue viruses, 348
cell lines, 348
Culicoides sonorensis, 348
field populations, 348
isozyme phenotyping, 348
Erethizon dorsatum,
carcass, 108, 108
carrion, 108, 108
decomposition, 108, 108
Dermacentor variabilis, 108, 108
Erethizon dorsatum, 108, 108
Eritrea,
Anopheles larval habitats, 921
larval control, 921
spatial temporal distribution, 921
esterase purification,
Boophilus microplus, 52
esterases, 52
insecticide resistance, 52

esterases,
Boophilus microplus, 52
esterase purification, 52
insecticide resistance, 52
Ethiopia,
Anopheles gambiae, 112
Anopheles quadriannulatus sp. B, 112
PCR assay, 112
Europe,
allozymes, 865
Anopheles claviger, 865
population genetics, 865
sibling species, 865
Evlachovaea,
Chagas' disease, 451
entomopathogenic fungi, 451
Triatoma sordida, 451
vector control, 451
excito-repellency,
Anopheles culicifacies, 58
bifenthrin, 58
malathion, 58
spraying, 58
vector density, 58
exophagy,
Anopheles minimus, 876
blood feeding, 876
mosquito abundance, 876
Thailand, 876
experimental infection,
antibody, 968
chronic infection, 968
St. Louis encephalitis virus, 968
western equine encephalomyelitis virus, 968

F

factors affecting immatures,
Ochlerotatus taeniorhynchus, 387
San Diego County, 387
Falconiformes,
Argas (*P.*) *keiransi* n. sp., 766
Chile, 766
fecundity,
Australia, 897
autogeny, 897
Ochlerotatus vigilax, 897
wing length, 897
feeding,
circadian activity, 903
female mosquito, 903
flow-through respirometry, 903
metabolic rate, 903
feeding by larval mosquitoes,
Bacillus thuringiensis israelensis, 860
maize pollen, 860
phagostimulant, 860
feeding success,
Borrelia bissettii, 311
Ixodes pacificus, 311
Ixodes spinipalpis, 311
rodent reservoir, 311
tick vector, 311
female mosquito,
circadian activity, 903
feeding, 903
flow-through respirometry, 903
metabolic rate, 903
field populations,
bluetongue viruses, 348
cell lines, 348
Culicoides sonorensis, 348
epizootic hemorrhagic disease viruses, 348
isozyme phenotyping, 348
field transmission,
arbovirus, 253
infection rate, 253
West Nile virus, 253
flagellin,
16S rRNA, 590
Amblyomma, 590
Borrelia, 590
Missouri, 590
fleas,
Deltadust, 718
epizootic, 718
insecticide, 718
plague, 718
prairie dogs, 718
Florida Keys,
Anopheles atropos, 361
Deinocerites cancer, 361
Ochlerotatus taeniorhynchus, 361
West Nile virus, 361
flow-through respirometry,
circadian activity, 903
feeding, 903
female mosquito, 903
metabolic rate, 903
forensic entomology, 259, 338, 559, 983
Anthropology Research Facility, 559
arthropod succession, 338
carrion ecology, 338
carrion-arthropod succession, 559
Chrysomya rufifacies, 259
Chrysomya villeneuvi, 983
hairy maggot, 983
larval morphology, 259
nutrient enrichment, 559
poaching, 338
scanning electron microscopy, 259
ultrastructure, 259
wildlife, 338

G

gene flow,
mitochondrial DNA, 30
Musca domestica, 30
single-strand conformation polymorphism, 30
genetic variability,
mosquitoes, 430
population structure, 430
random amplified polymorphic DNA, 430
genetic-relatedness malaria,
Anopheles dispersal, 421
kinship, 421
population structure, 421
genome sequence,
transgenics, 597
vector borne disease control, 597
geographic information systems (GIS),
Anopheles gambiae s.l. complex, 777
land use land cover change (LULC), 777
multispectral thermal imager (MTI), 777
geographic variation,
Culex pipiens quinquefasciatus, 882
isozyme electrophoresis, 882
population genetics, 882
temporal variation, 882
Getah,
isolation, 580
Japanese encephalitis, 580
Korea, 580
mosquitoes, 580
virus, 580
GIS,
environmental predictors, 570
Ixodes scapularis, 570
Poisson-normal model, 570
soil, 570
spatial correlation, 570
grasshopper,
bovine, 957
Melanoplus sanguinipes, 957
vesicular stomatitis virus, 957
VSV, 957
growth,
bacteria, 22
development, 22
Hematobia irritans, 22
larval survival, 22
media, 22

Guatemala,
Chagas, 436
population genetics, 436
Triatoma dimidiata, 436
vector control, 436

H

habitat,
adults, 18
Chagas' disease, 18
kissing bugs, 18
nymphs, 18
Haemaphysalis ornithophila,
Amblyomma testudinarium, 230
rickettsiae, 230
Thailand, 230
tick, 230
hairy maggot,
Chrysomya villeneuvi, 983
forensic entomology, 983
head lice, 468, 628, 795
Anoplura, 795
artificial feeding, 628
body louse, 468
insecticide-resistance, 468
in vitro rearing, 795
para-orthologous sodium channel gene, 468
pediculicide, 468
Pediculidae, 628, 795
Pediculus humanus capitis, 628, 795
Phthiraptera, 628
Hemaphysalis,
host, 103
Ixodes, 103
parasite, 103
phenology, 103
ticks, 103
hematin,
Ixodes scapularis, 849
pheromone, 849
purines, 849
Hematobia irritans,
bacteria, 22
development, 22
growth, 22
larval survival, 22
media, 22
hemocyte class,
ecdysial phase, 770
Ornithodoros moubata, 770
ultrastructure, 770
hemoglobin fragment,
antibacterial activity, 78
midgut, 78
Ornithodoros moubata, 78
highland malaria,
Anopheles spp., 706
malaria transmission, 706
Tanzania, 706
vectorial capacity, 706
Honduras,
Culex pseudes, 306
Mexico, 306
Ochlerotatus taeniorhynchus, 306
transmission, 306
Venezuelan equine encephalomyelitis virus, 306
horn fly,
diazinon, 942
insecticide resistance, 942
kdr, 942
permethrin, 942
host,
Hemaphysalis, 103
Ixodes, 103
parasite, 103
phenology, 103
ticks, 103
host–parasite associations,
Chiroptera, 813
ectoparasites, 813
Myodopsylla insignis, 813
Myotis lucifugus, 813
host-seeking,
arbovirus, 807
biting rates, 807
developmental substrates, 807
Oropouche virus, 807
house dust mites,
carpet cleaning, 508
Dermatophagoides, 508
Disodium octaborate tetrahydrate, 508
house finches, 206, 300
antibody, 300
chronic infection, 206
cyclophosphamide, 206
dexamethasone, 206
immunosuppression, 206
mosquito blood feeding, 206
St. Louis encephalitis virus, 300
viral persistence, 300
human bot fly,
Dermatobia hominis, 133
larva, 133
midgut, 133
ultrastructure, 133
hydrology,
amplification, 547
Culex nigripalpus, 547
St. Louis encephalitis, 547
wild birds, 547
2-(2-hydroxyethyl)-1-piperidinecarboxylic acid 1-methylpropyl ester, (1*S*, 2′*S*)-2-methylpiperidinyl-3-cyclohexen-1-carboxamide, 293
Aedes aegypti, 293
Anopheles stephensi, 293
N,*N*-diethyl-3-methylbenzamide, 293

I

I. ricinus,
B. burgdorferi, 690
reservoir, 690
small rodents, 690
urban forests, 690
immunocytochemistry,
Diptera, 755
dorsal septum, 755
neuropeptides, 755
immunosuppression,
chronic infection, 206
cyclophosphamide, 206
dexamethasone, 206
house finch, 206
mosquito blood feeding, 206
in vitro rearing,
Anoplura, 795
head lice, 795
Pediculidae, 795
Pediculus humanus capitis, 795
indoor climate,
Anopheles survival, 657
blood, 657
sugar, 657
vectorial capacity, 657
infection, 518, 686
Amblyomma americanum, 686
disease transmission, 686
Israel turkey meningoencepahlitis virus, 518
Rickettsia parkeri, 686
vector incrimination, 518
infection rate,
arbovirus, 253
field transmission, 253
West Nile virus, 253
infectivity,
Baculoviridae, 512
Culex, 512
CuniNPV, 512
mosquito, 512
infracommunities,
Acari, 996
bats, 996
Mexico, 996
Tadarida brasiliensis mexicana, 996
infusion,
amino acids, 441
Anopheles stephensi, 441

mating, 441
oogenesis, 441
inhibition,
deet, 463
dehydrolinalool, 463
linalool, 463
olfactometer, 463
insect growth regulator,
naturalyte, 475
neem, 475
storage mite, 475
Tyrophagus putrescentiae, 475
insect repellents,
Africanized honey bees, 275
Apis mellifera, 275
attacks, 275
citronella, 275
NN-diethyl-*m*-toluamide, 275
insecticide,
Deltadust, 718
epizootic, 718
fleas, 718
plague, 718
prairie dogs, 718
insecticide combinations,
Anopheles gambiae, 985
Côte d'Ivoire, 985
insecticide-treated nets, 985
insensitive acetylcholinesterase, 985
insecticide resistance, 52, 468, 491, 942
Anopheles gambiae, 491
body louse, 468
Boophilus microplus, 52
Cameroon, 491
diazinon, 942
esterase purification, 52
esterases, 52
head louse, 468
horn fly, 942
kdr, 942
para-orthologous sodium channel gene, 468
pediculicide, 468
permethrin, 942
insecticide-treated nets,
Anopheles gambiae, 985
Côte d'Ivoire, 985
insecticide combinations, 985
insensitive acetylcholinesterase, 985
insensitive acetylcholinesterase,
Anopheles gambiae, 985
Côte d'Ivoire, 985
insecticide combinations, 985
insecticide-treated nets, 985
interferon α/β,
mosquito, 199
saliva, 199
vesicular stomatitis virus, 199
introduction,
distributional range, 119
Ixodes persulcatus, 119
northern range border, 119
reproductive range, 119
Sakha Republic (Yakutia), 119
invasion,
archeological site, 800
Chagas disease, 800
sylvatic habitats, 800
Triatoma dimidiata, 800
isoenzymes,
morphology, 411
morphometry, 411
numerical taxonomy, 411
Panstrongylus megistus, 411
population, 411
isolation,
Getah, 580
Japanese encephalitis, 580
Korea, 580
mosquitoes, 580
virus, 580
isozyme electrophoresis,
Culex pipiens quinquefasciatus, 882
geographic variation, 882
population genetics, 882
temporal variation, 882
isozyme phenotyping,
bluetongue viruses, 348
cell lines, 348
Culicoides sonorensis, 348
epizootic hemorrhagic disease viruses, 348
field populations, 348
isozymes,
Lutzomyia longipalpis, 615
mitochondrial DNA, 615
phylogeny, 615
species complex, 615
Israel,
archaeology, 585
body louse, 585
Masada, 585
Pediculus humanus humanus, 585
Israel turkey meningoencepahlitis virus,
infection, 518
vector incrimination, 518
Ixodes, 103, 540
Birds, 540
Borrelia burgdorferi, 540
Hemaphysalis, 103
host, 103
Lyme disease, 540
parasite, 103
phenology, 103
risk factors, 540
ticks, 103
Ixodes angustus,
Alaska, 682
Leptinus occidentamericanus, 682
mammalian hosts, 682
Ixodes pacificus, 311, 534
Anaplasma phagocytophilum, 534
Borrelia bissettii, 311
Borrelia burgdorferi, 534
California, 534
Ehrlichia chaffeensis, 534
feeding success, 311
Ixodes spinipalpis, 311
rodent reservoir, 311
tick vector, 311
Ixodes persulcatus,
distributional range, 119
introduction, 119
northern range border, 119
reproductive range, 119
Sakha Republic (Yakutia), 119
Ixodes ricinus, 375, 723
biotopes, 375
blood meal analysis, 723
reservoir host identification, 723
Sweden, 375
tick sampling, 375
tick-borne zoonoses, 723
vegetation types, 375
Ixodes scapularis, 91, 179, 395, 528, 555, 570, 642, 732, 849, 964
Amblyomma americanum, 395, 528, 732
blacklegged tick, 91, 732
Borrelia burgdorferi, 91, 555, 964
deer density, 179
deer reduction, 642
Dermacentor andersoni, 528
Dermacentor variabilis, 528
distribution, 555
environmental predictors, 570
GIS, 570
hematin, 849
Ixodiphagus hookeri, 642
lone star tick, 732
Louisiana, 964
Lyme borreliosis, 964
Lyme disease, 91
Ontario, 91
parasitoid, 642
pheromone, 849
Poisson-normal model, 570
purines, 849
questing, 395
Rondeau Provincial Park, 91
soil, 570
spatial correlation, 570
tick, 642

weather, 395
West Nile virus, 528
white-tailed deer, 179
Ixodes schulzei,
Brazil, 116
laboratory rearing, 116
ticks, 116
Ixodes spinipalpis,
Borrelia bissettii, 311
feeding success, 311
Ixodes pacificus, 311
rodent reservoir, 311
tick vector, 311
Ixodiphagus hookeri,
deer reduction, 642
Ixodes scapularis, 642
parasitoid, 642
tick, 642

J

Japanese encephalitis, 82, 165, 249, 580
arbovirus, 249
Australia, 82, 249
Culex annulirostris, 165
Culex palpalis, 165
Culicidae, 165
dispersal, 165
Getah, 580
isolation, 580
Korea, 580
mosquito trap, 249
mosquitoes, 82, 580
polymerase chain reaction, 249
surveillance, 249
vector competence, 82
virus, 580

K

kairomones,
behavioral bioassay, 653
mosquito attractants, 653
KBR 3023,
Amblyomma hebraeum, 245
DEET, 245
piperidine, 245
repellents, 245
tick, 245
kdr,
diazinon, 942
insecticide resistance, 942
permethrin, 942
the horn fly, 942
kdr mutation,
Anopheles gambiae, 195
molecular M form, 195
tropical savannah area, 195
kinship,
Anopheles dispersal, 421
genetic-relatedness malaria, 421
population structure, 421
kissing bugs,
adults, 18
Chagas' disease, 18
habitat, 18
nymphs, 18
Korea,
Getah, 580
isolation, 580
Japanese encephalitis, 580
mosquitoes, 580
virus, 580

L

La Crosse virus,
Aedes albopictus, 403
arbovirus, 403
encephalitis, 403
Ochlerotatus triseriatus, 403
oviposition, 403
laboratory colony,
cold storage, 185
Dermacentor andersoni, 185
Rocky Mountain wood tick, 185
selection, 185
tick paralysis, 185
laboratory rearing,
Brazil, 116
Ixodes schulzei, 116
ticks, 116
land use land cover change (LULC),
Anopheles gambiae s.l. complex, 777
geographic information systems (GIS), 777
multispectral thermal imager (MTI), 777
larva,
Dermatobia hominis, 133
human bot fly, 133
midgut, 133
ultrastructure, 133
larval control,
Anopheles larval habitats, 921
Eritrea, 921
spatial temporal distribution, 921
larval morphology,
Chrysomya rufifacies, 259
forensic entomology, 259
scanning electron microscopy, 259
ultrastructure, 259
larval survival,
bacteria, 22
development, 22
growth, 22
Hematobia irritans, 22
media, 22
leishmaniasis, 141, 321
biting activity, 321
Paraphlebotomus, 141
Phlebotomus, 141, 321
resting sites, 321
Sand fly, 141
West Bank, 321
Leptinus occidentamericanus,
Alaska, 682
Ixodes angustus, 682
mammalian hosts, 682
Leptotrombidium,
chiggers, 64
personal protection, 64
repellents, 64
scrub typhus, 64
lethal ovitrap,
Aedes mosquitoes, 455
control, 455
dengue, 455
surveillance, 455
life tables,
Anopheles arabiensis, 737
seasonal conditions, 737
linalool,
deet, 463
dehydrolinalool, 463
inhibition, 463
olfactometer, 463
lone star tick,
Amblyomma americanum, 732
blacklegged tick, 732
Ixodes scapularis, 732
longevity,
Aedes albopictus, 522
mosquito strains, 522
vector competence, 522
Venezuelan equine encephalitis, 522
Louisiana,
Borrelia burgdorferi, 964
Ixodes scapularis, 964
Lyme borreliosis, 964
Lutzomyia,
Micropygomyia, 750
morphometrics, 750
multivariate analysis, 750
sand flies, 750
Lutzomyia longipalpis,
isozymes, 615
mitochondrial DNA, 615
phylogeny, 615
species complex, 615
Lyme borreliosis,
Borrelia burgdorferi, 964
Ixodes scapularis, 964
Louisiana, 964
Lyme disease, 91, 364, 540
Barbour-Stoenner-Kelly medium, 364
Birds, 540
blacklegged tick, 91

Borrelia burgdorferi, 91, 540
Ixodes, 540
Ixodes scapularis, 91
Ontario, 91
Peromyscus leucopus, 364
risk factors, 540
Rondeau Provincial Park, 91
spirochete, 364
tick, 364
lyme-like,
americanum, 100
B. lonestari, 100
STARI, 100

M

maize pollen,
Bacillus thuringiensis israelensis, 860
feeding by larval mosquitoes, 860
phagostimulant, 860
malaria, 68, 279, 636, 664
Amazon Region, 636
Anopheles, 636
Anopheles farauti, 68
Anopheles funestus, 664
bednet, 68
chromosomal inversions, 664
Papua New Guinea, 68
permethrin, 68
personal protection, 68
Plasmodium, 636
Sénégal, 279
species composition, 664
transmission, 279, 636
variability, 664
vectors, 279
malaria incidence,
Anopheles aquasalis, 379
Anopheles darlingi, 379
malaria vector, 379
northeastern Brazil, 379
malaria transmission,
Anopheles spp., 706
highland malaria, 706
Tanzania, 706
vectorial capacity, 706
malaria vector, 379, 991
Anopheles, 991
Anopheles aquasalis, 379
Anopheles darlingi, 379
malaria incidence, 379
northeastern Brazil, 379
trypsin, 991
malathion,
Anopheles culicifacies, 58
bifenthrin, 58
excito-repellency, 58
spraying, 58
vector density, 58
male mosquito,
autogeny, 150
mandible, 150
maxilla, 150
partial autogeny, 150
mammalian hosts,
Alaska, 682
Ixodes angustus, 682
Leptinus occidentamericanus, 682
mandible,
autogeny, 150
male mosquito, 150
maxilla, 150
partial autogeny, 150
mark-release,
Aedes aegypti, 6
Bootstrapping, 6
nonlinear regression, 6
statistical inference, 6
survival rates, 6
mark-release-recapture,
Culex tarsalis, 820
dispersal, 820
mosquitoes, 820
marsh deer,
Amblyomma cajennense, 268
Amblyomma triste, 268
Anocentor nitens, 268
Blastocerus dichotomus, 268
Boophilus microplus, 268
Masada,
archaeology, 585
body louse, 585
Israel, 585
Pediculus humanus humanus, 585
mating,
amino acids, 441
Anopheles stephensi, 441
infusion, 441
oogenesis, 441
maxilla,
autogeny, 150
male mosquito, 150
mandible, 150
partial autogeny, 150
media,
bacteria, 22
development, 22
growth, 22
Hematobia irritans, 22
larval survival, 22
Melanoplus sanguinipes,
bovine, 957
grasshopper, 957
vesicular stomatitis virus, 957
VSV, 957
metabolic rate,
circadian activity, 903
feeding, 903
female mosquito, 903
flow-through respirometry, 903
metamorphosis,
methoprene, 498
midgut, 498
mosquito, 498
methoprene,
metamorphosis, 498
midgut, 498
mosquito, 498
(1*S*, 2′*S*)-2-methylpiperidinyl-3-cyclohexen-1-carboxamide,
2-(2-hydroxyethyl)-1-piperidinecarboxylic acid 1-methylpropyl ester, 293
Aedes aegypti, 293
Anopheles stephensi, 293
N,N-diethyl-3-methylbenzamide, 293
Mexico, 306, 996
Acari, 996
bats, 996
Culex pseudes, 306
Honduras, 306
infracommunities, 996
Ochlerotatus taeniorhynchus, 306
Tadarida brasiliensis mexicana, 996
transmission, 306
Venezuelan equine encephalomyelitis virus, 306
Micropygomyia,
Lutzomyia, 750
morphometrics, 750
multivariate analysis, 750
sand flies, 750
midges,
abundance, 352
Culicoides, 352
parity, 352
midgut, 78, 133, 371, 498
Anopheles albimanus, 371
antibacterial activity, 78
bacteria, 371
Dermatobia hominis, 133
hemoglobin fragment, 78
human bot fly, 133
larva, 133
metamorphosis, 498
methoprene, 498
mosquito, 498
Ornithodoros moubata, 78
Plasmodium vivax, 371
sporogony, 371
ultrastructure, 133
midgut visceral muscles,
Aedes albopictus, 698
mosquito pathology, 698

salivary glands, 698
Sindbis virus, 698
Missouri,
16S rRNA, 590
Amblyomma, 590
Borrelia, 590
flagellin, 590
mite,
Acari, 146
Demodecidae, 146
domestic dog, 146
ectoparasite, 146
mitochondrial DNA, 30, 356, 615
Aedes albopictus, 356
gene flow, 30
isozymes, 615
Lutzomyia longipalpis, 615
Musca domestica, 30
phylogeny, 615
single-strand conformation polymorphism, 30
species complex, 615
Wolbachia, 356
wsp gene, 356
Mitonyssoides stercoralis,
bat guano habitat, 593
Coprolactistus whitakeri, 593
parasitism, 593
predation, 593
mixture,
Bacillus sphaericus, 672
Bacillus thuringiensis ssp. israelensis, 672
mosquitoes, 672
resistance, 672
model,
blacklegged tick, 238
weather, 238
white-tailed deer, 238
'4-poster', 238
molecular M form,
Anopheles gambiae, 195
kdr mutation, 195
tropical savannah area, 195
morphology,
isoenzymes, 411
morphometry, 411
numerical taxonomy, 411
Panstrongylus megistus, 411
population, 411
morphometrics,
Lutzomyia, 750
Micropygomyia, 750
multivariate analysis, 750
sand flies, 750
morphometry,
isoenzymes, 411
morphology, 411
numerical taxonomy, 411
Panstrongylus megistus, 411
population, 411
mosquito abundance,
Anopheles minimus, 876
blood feeding, 876
exophagy, 876
Thailand, 876
mosquito attractants,
behavioral bioassay, 653
kairomones, 653
mosquito blood feeding,
chronic infection, 206
cyclophosphamide, 206
dexamethasone, 206
house finch, 206
immunosuppression, 206
mosquito larval habitats,
Aedes albopictus, 73
Ochlerotatus triseriatus, 73
West Virginia, 73
mosquito pathology,
Aedes albopictus, 698
midgut visceral muscles, 698
salivary glands, 698
Sindbis virus, 698
mosquito strains,
Aedes albopictus, 522
longevity, 522
vector competence, 522
Venezuelan equine encephalitis, 522
mosquito trap,
arbovirus, 249
Australia, 249
Japanese encephalitis, 249
polymerase chain reaction, 249
surveillance, 249
mosquitoes, 1, 82, 199, 223, 430, 498, 512, 580, 672, 820
arbovirus, 223
Arkansas, 223
Australia, 82
Bacillus sphaericus, 672
Bacillus thuringiensis ssp. israelensis, 672
Baculoviridae, 512
blue jay, 223
Culex, 512
Culex tarsalis, 820
CuniNPV, 512
dispersal, 820
endosymbionts, 1
genetic variability, 430
Getah, 580
infectivity, 512
interferon α/β, 199
isolation, 580
Japanese encephalitis, 82, 580
Korea, 580
mark-release-recapture, 820
metamorphosis, 498
methoprene, 498
midgut, 498
mixture, 672
phylogeny, 1
population structure, 430
random amplified polymorphic DNA, 430
resistance, 672
saliva, 199
surveillance, 223
vector competence, 82
vesicular stomatitis virus, 199
virus, 580
West Nile virus, 223
Wolbachia, 1
wsp sequences, 1
multiplex polymerase chain reaction,
Bartonella, 329
black-tailed prairie dog, 329
Oropsylla, 329
Rickettsia, 329
Yersinia pestis, 329
multispectral thermal imager (MTI),
Anopheles gambiae s.l. complex, 777
geographic information systems (GIS), 777
land use land cover change (LULC), 777
multivariate analysis,
Lutzomyia, 750
Micropygomyia, 750
morphometrics, 750
sand flies, 750
Musca domestica,
gene flow, 30
mitochondrial DNA, 30
single-strand conformation polymorphism, 30
Myodopsylla insignis,
Chiroptera, 813
ectoparasites, 813
host–parasite associations, 813
Myotis lucifugus, 813
Myotis lucifugus,
Chiroptera, 813
ectoparasites, 813
host–parasite associations, 813
Myodopsylla insignis, 813

N

N,N-diethyl-3-methylbenzamide,
(1*S*, 2′*S*)-2-methylpiperidinyl-3-cyclohexen-1-carboxamide, 293
2-(2-hydroxyethyl)-1-piperidinecarboxylic acid 1-methylpropyl ester, 293
Aedes aegypti, 293
Anopheles stephensi, 293

N,N-diethyl-*m*-toluamide,
Africanized honey bees, 275
Apis mellifera, 275
attacks, 275
citronella, 275
insect repellents, 275
naturalyte,
insect growth regulator, 475
neem, 475
storage mite, 475
Tyrophagus putrescentiae, 475
neem,
insect growth regulator, 475
naturalyte, 475
storage mite, 475
Tyrophagus putrescentiae, 475
neuropeptides,
Diptera, 755
dorsal septum, 755
immunocytochemistry, 755
nitenpyram,
behavioral observations, 678
Capstar, 678
cat flea, 678
Ctenocephalides felis, 678
nonlinear regression,
Aedes aegypti, 6
Bootstrapping, 6
mark-release, 6
statistical inference, 6
survival rates, 6
non-native,
Arachnida, 159
range expansion, 159
Tegenaria agrestis, 159
northeastern Brazil, 379, 989
Anopheles aquasalis, 379
Anopheles darlingi, 379
Dasypodidae, 989
malaria incidence, 379
malaria vector, 379
Panstrongylus lutzi, 989
silvatic ecology, 989
Triatominae, 989
northern range border,
distributional range, 119
introduction, 119
Ixodes persulcatus, 119
reproductive range, 119
Sakha Republic (Yakutia), 119
numerical taxonomy,
isoenzymes, 411
morphology, 411
morphometry, 411
Panstrongylus megistus, 411
population, 411
nutrient enrichment,
Anthropology Research Facility, 559
carrion-arthropod succession, 559
forensic entomology, 559
nymphs,
adults, 18
Chagas' disease, 18
habitat, 18
kissing bugs, 18

O

occurrence,
abundance, 785
Brazil, 785
dengue vectors, 785
United States, 785
Ochlerotatus caspius,
area repellent, 930
cis/*trans* allethrin, 930
Phlebotomus papatasi, 930
Thermacell, 930
Ochlerotatus taeniorhynchus, 306, 361, 387
Anopheles atropos, 361
Culex pseudes, 306
Deinocerites cancer, 361
factors affecting immatures, 387
Florida Keys, 361
Honduras, 306
Mexico, 306
San Diego County, 387
transmission, 306
Venezuelan equine encephalomyelitis virus, 306
West Nile virus, 361
Ochlerotatus triseriatus, 73, 403
Aedes albopictus, 73, 403
arbovirus, 403
encephalitis, 403
La Crosse virus, 403
mosquito larval habitats, 73
oviposition, 403
West Virginia, 73
Ochlerotatus vigilax,
Australia, 897
autogeny, 897
fecundity, 897
wing length, 897
olfaction,
attractants, 284
carbon dioxide, 284
Culicoides furens, 284
electrophysiology, 284
olfactometer,
deet, 463
dehydrolinalool, 463
inhibition, 463
linalool, 463
Ontario,
blacklegged tick, 91
Borrelia burgdorferi, 91
Ixodes scapularis, 91
Lyme disease, 91
Rondeau Provincial Park, 91
oogenesis,
amino acids, 441
Anopheles stephensi, 441
infusion, 441
mating, 441
organic infusion,
Aedes albopictus, 841
attractants, 841
bacteria, 841
electroantennogram, 841
oviposition, 841
organophosphate,
acaricide resistance, 482
Boophilus microplus, 482
cattle tick, 482
coumaphos, 482
diazinon, 482
orientation,
Aedes vexans, 607
Anopheles walkeri, 607
carbon dioxide, 607
Coquillettidae perturbans, 607
Ornithodoros moubata, 78, 770
antibacterial activity, 78
ecdysial phase, 770
hemocyte class, 770
hemoglobin fragment, 78
midgut, 78
ultrastructure, 770
Oropouche virus,
arbovirus, 807
biting rates, 807
developmental substrates, 807
host-seeking, 807
Oropsylla,
Bartonella, 329
black-tailed prairie dog, 329
multiplex polymerase chain reaction, 329
Rickettsia, 329
Yersinia pestis, 329
oviposition, 403, 830, 841
Aedes aegypti, 830
Aedes albopictus, 403, 841
arbovirus, 403
attractants, 841
bacteria, 841
biological control, 830
electroantennogram, 841
encephalitis, 403
La Crosse virus, 403
Ochlerotatus triseriatus, 403
organic infusion, 841
Plagiorchis elegans, 830
population ecology, 830

P

Panstrongylus lutzi,
Dasypodidae, 989
northeastern Brazil, 989
silvatic ecology, 989
Triatominae, 989

Panstrongylus megistus,
isoenzymes, 411
morphology, 411
morphometry, 411
numerical taxonomy, 411
population, 411
Papua New Guinea,
Anopheles farauti, 68
bednet, 68
malaria, 68
permethrin, 68
personal protection, 68
Parafilm M,
Aedes, 588
artificial feeding, 588
para-orthologous sodium channel gene,
body louse, 468
head louse, 468
insecticide-resistance, 468
pediculicide, 468
Paraphlebotomus,
leishmaniasis, 141
Phlebotomus, 141
Sand fly, 141
parasite,
Hemaphysalis, 103
host, 103
Ixodes, 103
phenology, 103
ticks, 103
parasitism,
bat guano habitat, 593
Coprolactistus whitakeri, 593
Mitonyssoides stercoralis, 593
predation, 593
parasitoid,
deer reduction, 642
Ixodes scapularis, 642
Ixodiphagus hookeri, 642
tick, 642
parity,
abundance, 352
Culicoides, 352
midges, 352
partial autogeny,
autogeny, 150
male mosquito, 150
mandible, 150
maxilla, 150
PCR assay,
Anopheles gambiae, 112
Anopheles quadriannulatus sp. B, 112
Ethiopia, 112
pediculicide,
body louse, 468
head louse, 468
insecticide-resistance, 468
para-orthologous sodium channel gene, 468
Pediculidae, 628, 795
Anoplura, 795
artificial feeding, 628
head lice, 628, 795
in vitro rearing, 795
Pediculus humanus capitis, 628, 795
Phthiraptera, 628
Pediculus humanus capitis, 447, 628, 795
Anoplura, 795
artificial feeding, 628
bioassay, 447
head lice, 628, 795
in vitro rearing, 795
Pediculidae, 628, 795
permethrin resistance, 447
Phthiraptera, 628
Pediculus humanus humanus,
archaeology, 585
body louse, 585
Israel, 585
Masada, 585
permethrin, 68, 935, 942
Anopheles farauti, 68
arthropods, 935
bednet, 68
diazinon, 942
insecticide resistance, 942
kdr, 942
malaria, 68
Papua New Guinea, 68
personal protection, 68
repellents, 935
the horn fly, 942
uniforms, 935
permethrin resistance,
bioassay, 447
Pediculus humanus capitis, 447
Peromyscus leucopus,
Barbour-Stoenner-Kelly medium, 364
Lyme disease, 364
spirochete, 364
tick, 364
personal protection, 64, 68
Anopheles farauti, 68
bednet, 68
chiggers, 64
Leptotrombidium, 64
malaria, 68
Papua New Guinea, 68
permethrin, 68
repellents, 64
scrub typhus, 64
phagostimulant,
Bacillus thuringiensis israelensis, 860
feeding by larval mosquitoes, 860
maize pollen, 860
phenology,
Hemaphysalis, 103
host, 103
Ixodes, 103
parasite, 103
ticks, 103
pheromone,
hematin, 849
Ixodes scapularis, 849
purines, 849
Phlebotomus, 141, 321
biting activity, 321
leishmaniasis, 141, 321
Paraphlebotomus, 141
resting sites, 321
Sand fly, 141
West Bank, 321
Phlebotomus papatasi,
area repellent, 930
cis/*trans* allethrin, 930
Ochlerotatus caspius, 930
Thermacell, 930
Phthiraptera,
artificial feeding, 628
head lice, 628
Pediculidae, 628
Pediculus humanus capitis, 628
phylogeny, 1, 615
endosymbionts, 1
isozymes, 615
Lutzomyia longipalpis, 615
mitochondrial DNA, 615
mosquitoes, 1
species complex, 615
Wolbachia, 1
wsp sequences, 1
piperidine,
Amblyomma hebraeum, 245
DEET, 245
KBR 3023, 245
repellents, 245
tick, 245
Plagiorchis elegans,
Aedes aegypti, 830
biological control, 830
oviposition, 830
population ecology, 830
plague,
Deltadust, 718
epizootic, 718
fleas, 718
insecticide, 718
prairie dogs, 718
Plasmodium,
Amazon Region, 636
Anopheles, 636
malaria, 636
transmission, 636
Plasmodium vivax,
Anopheles albimanus, 371
bacteria, 371

midgut, 371
sporogony, 371
poaching,
arthropod succession, 338
carrion ecology, 338
forensic entomology, 338
wildlife, 338
Poisson-normal model,
environmental predictors, 570
GIS, 570
Ixodes scapularis, 570
soil, 570
spatial correlation, 570
polymerase chain reaction,
arbovirus, 249
Australia, 249
Japanese encephalitis, 249
mosquito trap, 249
surveillance, 249
popcorn,
age-structure, 125
Culex pipiens, 125
vector-borne disease, 125
Wolbachia, 125
population,
isoenzymes, 411
morphology, 411
morphometry, 411
numerical taxonomy, 411
Panstrongylus megistus, 411
population ecology,
Aedes aegypti, 830
biological control, 830
oviposition, 830
Plagiorchis elegans, 830
population genetics, 436, 865, 882
allozymes, 865
Anopheles claviger, 865
Chagas, 436
Culex pipiens quinquefasciatus, 882
Europe, 865
geographic variation, 882
Guatemala, 436
isozyme electrophoresis, 882
sibling species, 865
temporal variation, 882
Triatoma dimidiata, 436
vector control, 436
population structure, 421, 430
Anopheles dispersal, 421
genetic variability, 430
genetic-relatedness malaria, 421
kinship, 421
mosquitoes, 430
random amplified polymorphic DNA, 430
'4-poster',
blacklegged tick, 238
model, 238
weather, 238
white-tailed deer, 238
prairie dogs,
Deltadust, 718
epizootic, 718
fleas, 718
insecticide, 718
plague, 718
predation,
bat guano habitat, 593
Coprolactistus whitakeri, 593
Mitonyssoides stercoralis, 593
parasitism, 593
premolt period,
Amblyomma cajennense, 170
Brazil, 170
diapause, 170
purines,
hematin, 849
Ixodes scapularis, 849
pheromone, 849
pyrethroid insecticides,
domestic control, 912
Triatoma barberi, 912
Triatoma pallidipennis, 912

Q

questing,
Amblyomma americanum, 395
Ixodes scapularis, 395
weather, 395

R

random amplified polymorphic DNA,
genetic variability, 430
mosquitoes, 430
population structure, 430
range expansion,
Arachnida, 159
non-native, 159
Tegenaria agrestis, 159
rapid detection,
arbovirus, 95
St. Louis encephalitis, 95
surveillance, 95
West Nile virus, 95
wicking assay, 95
repellents, 64, 245, 935
Amblyomma hebraeum, 245
arthropods, 935
chiggers, 64
DEET, 245
KBR 3023, 245
Leptotrombidium, 64
permethrin, 935
personal protection, 64
piperidine, 245
scrub typhus, 64
tick, 245
uniforms, 935
reproductive range,
distributional range, 119
introduction, 119
Ixodes persulcatus, 119
northern range border, 119
Sakha Republic (Yakutia), 119
reservoir,
B. burgdorferi, 690
I. ricinus, 690
small rodents, 690
urban forests, 690
reservoir host identification,
blood meal analysis, 723
Ixodes ricinus, 723
tick-borne zoonoses, 723
resistance,
Bacillus sphaericus, 672
Bacillus thuringiensis ssp. israelensis, 672
mixture, 672
mosquitoes, 672
resting sites,
biting activity, 321
leishmaniasis, 321
Phlebotomus, 321
West Bank, 321
Rhipicephalus sanguineus,
acetylcholinesterase, 890
cDNA, 890
Dermacentor variabilis, 890
ticks, 890
Rickettsia,
Bartonella, 329
black-tailed prairie dog, 329
multiplex polymerase chain reaction, 329
Oropsylla, 329
Yersinia pestis, 329
Rickettsia parkeri,
Amblyomma americanum, 686
disease transmission, 686
infection, 686
rickettsiae,
Amblyomma testudinarium, 230
Haemaphysalis ornithophila, 230
Thailand, 230
tick, 230
risk factors,
Birds, 540
Borrelia burgdorferi, 540
Ixodes, 540
Lyme disease, 540
Rocky Mountain wood tick,
cold storage, 185
Dermacentor andersoni, 185
laboratory colony, 185
selection, 185
tick paralysis, 185

rodent reservoir,
Borrelia bissettii, 311
feeding success, 311
Ixodes pacificus, 311
Ixodes spinipalpis, 311
tick vector, 311
Rondeau Provincial Park,
blacklegged tick, 91
Borrelia burgdorferi, 91
Ixodes scapularis, 91
Lyme disease, 91
Ontario, 91
16S rRNA,
Amblyomma, 590
Borrelia, 590
flagellin, 590
Missouri, 590

S

Sakha Republic (Yakutia),
distributional range, 119
introduction, 119
Ixodes persulcatus, 119
northern range border, 119
reproductive range, 119
saliva,
interferon α/β, 199
mosquito, 199
vesicular stomatitis virus, 199
salivary glands,
Aedes albopictus, 698
midgut visceral muscles, 698
mosquito pathology, 698
Sindbis virus, 698
San Diego County,
factors affecting immatures, 387
Ochlerotatus taeniorhynchus, 387
sand flies, 141, 750
leishmaniasis, 141
Lutzomyia, 750
Micropygomyia, 750
morphometrics, 750
multivariate analysis, 750
Paraphlebotomus, 141
Phlebotomus, 141
scanning electron microscopy,
Chrysomya rufifacies, 259
forensic entomology, 259
larval morphology, 259
ultrastructure, 259
scrub typhus,
chiggers, 64
Leptotrombidium, 64
personal protection, 64
repellents, 64
seasonal conditions,
Anopheles arabiensis, 737
life tables, 737
selection,
cold storage, 185
Dermacentor andersoni, 185
laboratory colony, 185
Rocky Mountain wood tick, 185
tick paralysis, 185
Sénégal,
malaria, 279
transmission, 279
vectors, 279
sibling species,
allozymes, 865
Anopheles claviger, 865
Europe, 865
population genetics, 865
silvatic ecology,
Dasypodidae, 989
northeastern Brazil, 989
Panstrongylus lutzi, 989
Triatominae, 989
Sindbis virus,
Aedes albopictus, 698
midgut visceral muscles, 698
mosquito pathology, 698
salivary glands, 698
single-strand conformation polymorphism,
gene flow, 30
mitochondrial DNA, 30
Musca domestica, 30
small rodents,
B. burgdorferi, 690
I. ricinus, 690
reservoir, 690
urban forests, 690
soil,
environmental predictors, 570
GIS, 570
Ixodes scapularis, 570
Poisson-normal model, 570
spatial correlation, 570
spatial correlation,
environmental predictors, 570
GIS, 570
Ixodes scapularis, 570
Poisson-normal model, 570
soil, 570
spatial temporal distribution,
Anopheles larval habitats, 921
Eritrea, 921
larval control, 921
species complex,
isozymes, 615
Lutzomyia longipalpis, 615
mitochondrial DNA, 615
phylogeny, 615
species composition,
Anopheles funestus, 664
chromosomal inversions, 664
malaria, 664
variability, 664
spiracular index,
Anopheles stephensi type form, 747
ecological variants, 747
taxonomic tools, 747
thoracic spiracle length, 747
variant *mysorensis*, 747
spirochete,
Barbour-Stoenner-Kelly medium, 364
Lyme disease, 364
Peromyscus leucopus, 364
tick, 364
sporogony,
Anopheles albimanus, 371
bacteria, 371
midgut, 371
Plasmodium vivax, 371
spraying,
Anopheles culicifacies, 58
bifenthrin, 58
excito-repellency, 58
malathion, 58
vector density, 58
St. Louis encephalitis, 95, 547
amplification, 547
arbovirus, 95
Culex nigripalpus, 547
hydrology, 547
rapid detection, 95
surveillance, 95
West Nile virus, 95
wicking assay, 95
wild birds, 547
St. Louis encephalitis virus, 300, 968
antibody, 300, 968
chronic infection, 968
experimental infection, 968
house finches, 300
viral persistence, 300
western equine encephalomyelitis virus, 968
STARI,
americanum, 100
B. lonestari, 100
lyme-like, 100
statistical inference,
Aedes aegypti, 6
Bootstrapping, 6
mark-release, 6
nonlinear regression, 6
survival rates, 6
storage mite,
insect growth regulator, 475
naturalyte, 475
neem, 475
Tyrophagus putrescentiae, 475
sugar,
Anopheles survival, 657
blood, 657

indoor climate, 657
vectorial capacity, 657
surveillance, 95, 223, 249, 455
Aedes mosquitoes, 455
arbovirus, 95, 223, 249
Arkansas, 223
Australia, 249
blue jay, 223
control, 455
dengue, 455
Japanese encephalitis, 249
lethal ovitrap, 455
mosquito, 223
mosquito trap, 249
polymerase chain reaction, 249
rapid detection, 95
St. Louis encephalitis, 95
West Nile virus, 95, 223
wicking assay, 95
survival rates,
Aedes aegypti, 6
Bootstrapping, 6
mark-release, 6
nonlinear regression, 6
statistical inference, 6
Sweden,
biotopes, 375
Ixodes ricinus, 375
tick sampling, 375
vegetation types, 375
sylvatic habitats,
archeological site, 800
Chagas disease, 800
invasion, 800
Triatoma dimidiata, 800

T

Tadarida brasiliensis mexicana,
Acari, 996
bats, 996
infracommunities, 996
Mexico, 996
Tanzania,
Anopheles spp., 706
highland malaria, 706
malaria transmission, 706
vectorial capacity, 706
taxonomic tools,
Anopheles stephensi type form, 747
ecological variants, 747
spiracular index, 747
thoracic spiracle length, 747
variant *mysorensis*, 747
Tegenaria agrestis,
Arachnida, 159
non-native, 159
range expansion, 159
temporal variation,
Culex pipiens quinquefasciatus, 882
geographic variation, 882
isozyme electrophoresis, 882
population genetics, 882
Thailand, 230, 876
Amblyomma testudinarium, 230
Anopheles minimus, 876
blood feeding, 876
exophagy, 876
Haemaphysalis ornithophila, 230
mosquito abundance, 876
rickettsiae, 230
tick, 230
Thermacell,
area repellent, 930
cis/*trans* allethrin, 930
Ochlerotatus caspius, 930
Phlebotomus papatasi, 930
thoracic spiracle length,
Anopheles stephensi type form, 747
ecological variants, 747
spiracular index, 747
taxonomic tools, 747
variant *mysorensis*, 747
tick paralysis,
cold storage, 185
Dermacentor andersoni, 185
laboratory colony, 185
Rocky Mountain wood tick, 185
selection, 185
tick sampling,
biotopes, 375
Ixodes ricinus, 375
Sweden, 375
vegetation types, 375
tick vector,
Borrelia bissettii, 311
feeding success, 311
Ixodes pacificus, 311
Ixodes spinipalpis, 311
rodent reservoir, 311
tick-borne zoonoses,
blood meal analysis, 723
Ixodes ricinus, 723
reservoir host identification, 723
ticks, 103, 116, 230, 245, 364, 642, 890
acetylcholinesterase, 890
Amblyomma hebraeum, 245
Amblyomma testudinarium, 230
Barbour-Stoenner-Kelly medium, 364
Brazil, 116
cDNA, 890
deer reduction, 642
DEET, 245
Dermacentor variabilis, 890
Haemaphysalis ornithophila, 230
Hemaphysalis, 103
host, 103
Ixodes, 103
Ixodes scapularis, 642
Ixodes schulzei, 116
Ixodiphagus hookeri, 642
KBR 3023, 245
laboratory rearing, 116
Lyme disease, 364
parasite, 103
parasitoid, 642
Peromyscus leucopus, 364
phenology, 103
piperidine, 245
repellents, 245
Rhipicephalus sanguineus, 890
rickettsiae, 230
spirochete, 364
Thailand, 230
transgenics,
genome sequence, 597
vector borne disease control, 597
transmission, 279, 306, 636, 950
Aedes aegypti, 950
Amazon Region, 636
Anopheles, 636
Culex pseudes, 306
dengue, 950
Honduras, 306
malaria, 279, 636
Mexico, 306
Ochlerotatus taeniorhynchus, 306
Plasmodium, 636
Sénégal, 279
vector competence, 950
vectors, 279
Venezuelan equine encephalomyelitis virus, 306
virus, 950
Triatoma barberi,
domestic control, 912
pyrethroid insecticides, 912
Triatoma pallidipennis, 912
Triatoma dimidiata, 436, 800
archeological site, 800
Chagas, 436
Chagas disease, 800
Guatemala, 436
invasion, 800
population genetics, 436
sylvatic habitats, 800
vector control, 436
Triatoma pallidipennis,
domestic control, 912
pyrethroid insecticides, 912
Triatoma barberi, 912

Triatoma sordida,
Chagas' disease, 451
entomopathogenic fungi, 451
Evlachovaea, 451
vector control, 451
Triatominae,
Dasypodidae, 989
northeastern Brazil, 989
Panstrongylus lutzi, 989
silvatic ecology, 989
tropical savannah area,
Anopheles gambiae, 195
kdr mutation, 195
molecular M form, 195
trypsin,
Anopheles, 991
malaria vector, 991
Tyrophagus putrescentiae,
insect growth regulator, 475
naturalyte, 475
neem, 475
storage mite, 475

U

ultrastructure, 133, 259, 770
Chrysomya rufifacies, 259
Dermatobia hominis, 133
ecdysial phase, 770
forensic entomology, 259
hemocyte class, 770
human bot fly, 133
larva, 133
larval morphology, 259
midgut, 133
Ornithodoros moubata, 770
scanning electron microscopy, 259
uniforms,
arthropods, 935
permethrin, 935
repellents, 935
United States,
abundance, 785
Brazil, 785
dengue vectors, 785
occurrence, 785
urban forests,
B. burgdorferi, 690
I. ricinus, 690
reservoir, 690
small rodents, 690

V

variability,
Anopheles funestus, 664
chromosomal inversions, 664
malaria, 664
species composition, 664
variant *mysorensis*,
Anopheles stephensi type form, 747
ecological variants, 747
spiracular index, 747
taxonomic tools, 747
thoracic spiracle length, 747
vector borne disease control,
genome sequence, 597
transgenics, 597
vector competence, 82, 522, 950
Aedes aegypti, 950
Aedes albopictus, 522
Australia, 82
dengue, 950
Japanese encephalitis, 82
longevity, 522
mosquito strains, 522
mosquitoes, 82
transmission, 950
Venezuelan equine encephalitis, 522
virus, 950
vector control, 436, 451
Chagas, 436
Chagas' disease, 451
entomopathogenic fungi, 451
Evlachovaea, 451
Guatemala, 436
population genetics, 436
Triatoma dimidiata, 436
Triatoma sordida, 451
vector density,
Anopheles culicifacies, 58
bifenthrin, 58
excito-repellency, 58
malathion, 58
spraying, 58
vector incrimination,
infection, 518
Israel turkey meningo-encepahlitis virus, 518
vector-borne disease,
age-structure, 125
Culex pipiens, 125
popcorn, 125
Wolbachia, 125
vectorial capacity, 657, 706
Anopheles spp., 706
Anopheles survival, 657
blood, 657
highland malaria, 706
indoor climate, 657
malaria transmission, 706
sugar, 657
Tanzania, 706
vectors, 279, 577
bluetongue, 577
Culicoides mohave, 577
Culicoides sonorensis, 577
epizootic hemorrhagic disease, 577
malaria, 279
Sénégal, 279
transmission, 279
vegetation types,
biotopes, 375
Ixodes ricinus, 375
Sweden, 375
tick sampling, 375
Venezuelan equine encephalitis,
Aedes albopictus, 522
longevity, 522
mosquito strains, 522
vector competence, 522
Venezuelan equine encephalomyelitis virus,
Culex pseudes, 306
Honduras, 306
Mexico, 306
Ochlerotatus taeniorhynchus, 306
transmission, 306
vertical transmission,
Culex pipiens, 743
Culex tarsalis, 743
West Nile virus, 743
vesicular stomatitis virus, 199, 957
bovine, 957
grasshopper, 957
interferon α/β, 199
Melanoplus sanguinipes, 957
mosquito, 199
saliva, 199
VSV, 957
viral persistence,
antibody, 300
house finches, 300
St. Louis encephalitis virus, 300
virus, 580, 950
Aedes aegypti, 950
dengue, 950
Getah, 580
isolation, 580
Japanese encephalitis, 580
Korea, 580
mosquitoes, 580
transmission, 950
vector competence, 950
VSV,
bovine, 957
grasshopper, 957
Melanoplus sanguinipes, 957
vesicular stomatitis virus, 957

W

weather, 238, 395
Amblyomma americanum, 395
blacklegged tick, 238
Ixodes scapularis, 395
model, 238
questing, 395
white-tailed deer, 238
'4-poster', 238

West Bank,
biting activity, 321
leishmaniasis, 321
Phlebotomus, 321
resting sites, 321
West Nile,
bloodmeal, 215
Culex, 215
West Nile virus, 95, 223, 253, 361, 528, 743
Amblyomma americanum, 528
Anopheles atropos, 361
arbovirus, 95, 223, 253
Arkansas, 223
blue jay, 223
Culex pipiens, 743
Culex tarsalis, 743
Deinocerites cancer, 361
Dermacentor andersoni, 528
Dermacentor variabilis, 528
field transmission, 253
Florida Keys, 361
infection rate, 253
Ixodes scapularis, 528
mosquito, 223
Ochlerotatus taeniorhynchus, 361
rapid detection, 95
St. Louis encephalitis, 95
surveillance, 95, 223
vertical transmission, 743
wicking assay, 95
West Virginia,
Aedes albopictus, 73
mosquito larval habitats, 73
Ochlerotatus triseriatus, 73
western equine encephalomyelitis virus,
antibody, 968
chronic infection, 968
experimental infection, 968
St. Louis encephalitis virus, 968
white-tailed deer, 179, 238
blacklegged tick, 238
deer density, 179
Ixodes scapularis, 179
model, 238
weather, 238
'4-poster', 238
wicking assay,
arbovirus, 95
rapid detection, 95
St. Louis encephalitis, 95
surveillance, 95
West Nile virus, 95
wild birds,
amplification, 547
Culex nigripalpus, 547
hydrology, 547
St. Louis encephalitis, 547
wildlife,
arthropod succession, 338
carrion ecology, 338
forensic entomology, 338
poaching, 338
wing length,
Australia, 897
autogeny, 897
fecundity, 897
Ochlerotatus vigilax, 897
Wolbachia, 1, 125, 356
Aedes albopictus, 356
age-structure, 125
Culex pipiens, 125
endosymbionts, 1
mitochondrial DNA, 356
mosquitoes, 1
phylogeny, 1
popcorn, 125
vector-borne disease, 125
wsp gene, 356
wsp sequences, 1
wsp gene,
Aedes albopictus, 356
mitochondrial DNA, 356
Wolbachia, 356
wsp sequences,
endosymbionts, 1
mosquitoes, 1
phylogeny, 1
Wolbachia, 1

Y

Yersinia pestis,
Bartonella, 329
black-tailed prairie dog, 329
multiplex polymerase chain reaction, 329
Oropsylla, 329
Rickettsia, 329